



**EUREX** *Medica*®

**EUREX MEDICA, spol. s r.o.**

Výstavní 604/111

703 00 Ostrava – Vítkovice

Tel.: 599 526 510,

Fax: 596 614 507

e-mail: [expedice@eurexmedica.cz](mailto:expedice@eurexmedica.cz)

web: <http://www.eurexmedica.cz>



# Product list

Your trusted partner in diagnostics

Blood grouping serology | Bacteriological test reagents | Culture media | Immunoassays  
Monoclonal antibodies | Contract manufacturing | MICRONAUT system

# Products

DIAGNOSTICS WITH PASSION



**sifin diagnostics gmbh**  
Berliner Allee 317-321  
13088 Berlin, Germany

Phone: +49 30 700 144-0  
Telefax: +49 30 700 144-30  
E-Mail: [info@sifin.de](mailto:info@sifin.de)



**EUREX Medica**<sup>®</sup>

**EUREX MEDICA, spol. s r.o.**

Výstavní 604/111

703 00 Ostrava – Vítkovice

Tel.: 599 526 510,

Fax: 596 614 507

e-mail: [expedice@eurexmedica.cz](mailto:expedice@eurexmedica.cz)

web: <http://www.eurexmedica.cz>

# Contents

## Miscellaneous

Contact .....	2
Company profile .....	3
Order blank .....	4

## Blood Grouping Serology

Bedside cards for identity proof .....	6
Antigen determination of the ABO system .....	6
Antigen determination of the Rh system .....	6
Detection of additional blood grouping antigens .....	7
Anti-Human Globulin (Coombs) .....	7
Enzymes .....	7
Kits for PK7200® and PK7300® .....	8
Test reagents for PK7200® and PK7300® .....	9
Monoclonal antibodies .....	10

## Bacteriological Test Reagents

Salmonella diagnostics .....	12
Shigella diagnostics .....	20
Yersinia diagnostics .....	22
Coli diagnostics .....	23

## Culture Media

Dehydrated culture media .....	28
Supplements .....	42
Additives .....	44
Detection reagents .....	44
Ready-to-use culture media .....	45
Ready-to-use culture media for breweries .....	45
Base materials .....	46
Index, alphabetical .....	62

Certificate DIN EN ISO 9001 .....	68
Certificate DIN EN ISO 13485 .....	69
Certificate DIN EN ISO/IEC 17025 .....	70
Distributors .....	71
Index .....	62

## Immunoassays

Detection of rabies virus .....	47
---------------------------------	----

## Monoclonal Antibodies

Contract manufacturing monoclonal antibodies .....	49
Detection of human IgM antibodies .....	50
Anti-Mistletoe lectins .....	50
Shigatoxin (Verotoxin) specific monoclonal antibodies .....	50
Salmonella specific antibodies .....	50

## Contract Manufacturing

Contract manufacturing .....	51
------------------------------	----

## MICRONAUT System

MICRONAUT identification .....	54
MICRONAUT antimicrobial susceptibility test (AST) .....	55
Customer defined plates .....	55
Standard plates for clinical laboratories .....	55
Standard plates for veterinary laboratories .....	56
Standard plates for antifungal susceptibility testing of yeasts .....	56
MICRONAUT MIC Strips .....	56
Special plates for clinical laboratories .....	57
Special plates for interlaboratory comparison .....	58
MICRONAUT reagents .....	58
MICRONAUT media .....	59
MICRONAUT software .....	60
Instruments .....	60
Consumables .....	61

# Your contact with sifin

## You may reach us personally German time

Monday - Thursday

7:30 am to 4:15 pm

Friday

7:30 am to 3:00 pm

Head office

Telephone: +49 30 700 144-0

Telefax: +49 30 700 144-30

## Export department

**Uwe Stoehr**

Telephone: +49 30 700 144-285

Telefax: +49 30 700 144-30

E-Mail: [u.stoehr@sifin.de](mailto:u.stoehr@sifin.de)

**Ronald Rasche**

Telephone: +49 30 700 144-222

Telefax: +49 30 700 144-30

E-Mail: [r.rasche@sifin.de](mailto:r.rasche@sifin.de)

**Mario Konojacki**

Telephone: +49 30 700 144-283

Telefax: +49 30 700 144-30

E-Mail: [m.konojacki@sifin.de](mailto:m.konojacki@sifin.de)

## Acceptance of order

Telephone: +49 30 700 144-222 or

+49 30 700 144-283

Telefax: +49 30 700 144-30

E-Mail: [info@sifin.de](mailto:info@sifin.de)

# Company Profile

DIAGNOSTICS WITH PASSION



## **Your Trusted Partner in Diagnostics**

sifin diagnostics gmbh has emerged from the former 'Staatliches Institut für Immunpräparate und Nährmedien' of the GDR. Thanks to investments by a German family of entrepreneurs in 1992, the diagnostics division (microbiology and blood grouping serology) survived after the reunification.

Since then, sifin diagnostics gmbh has recorded constant growth. The number of employees and turnover have tripled, and today the company, which is still in family ownership, has an established

position in the diagnostics industry. In the past few years, especially the export activities have been strengthened and could be intensively developed in the key markets of Europe and Asia.

**sifin diagnostics gmbh offers more than 600 standard products in the area of microbiology and immunology. Beside this we are proud of our flexibility in special requests. We thus provide culture media according to your formulation, or we take care of your cell line. Our monoclonal antibodies may be supplied in your required format as ready-to-use or concentrate in bulk volumes.**

The company has an excellent reputation thanks to its great flexibility and reliability in the fulfilment of specific customer requests. sifin diagnostics gmbh is a trusted supplier and competent contact for many companies all over the world.





# Blood Grouping Serology

DIAGNOSTICS WITH PASSION



## Blood grouping for every format

Our experience in development and production of monoclonal antibodies for blood grouping has a long history. For more than 40 years we have been working with monoclonal antibodies for blood grouping.

Our products are available in different ready-to-use formats: dropper bottle for routine diagnostics, bedside cards for confirmation tests or reagents for PK® instruments.

To IVD companies we offer our monoclonal antibodies in high concentration or in bulk.

## Our portfolio comprises:

- Bedside cards
- Ready-to-use reagents
- Ready-to-use reagents as bulk and concentrates
- Ready-to-use test reagents for blood grouping on PK® instruments



# Blood Grouping Serology

## Bedside cards for identity proof

Ready-to-use twin-cards Serafol® AB0 and Serafol® AB0+D are used for an identity check immediately prior to a blood transfusion (bedside test). The test is the confirmation of earlier ABO and D blood typing of the recipient and ensures the compatibility of blood types of the recipient and the blood to be transfused. Thus, possible mismatches can be detected. Cards are coated with dried monoclonal reagents. The test principle is a hemagglutination test for the detection of the respective red cell antigens. ABO blood types and the rhesus characteristic D are defined by the presence or absence of the antigens A, B, and D on red cells. If such antigens are present on the red cells, they will be agglutinated by the corresponding antibodies (positive reaction).

Art. No.	Product	Description	Packing
BG1721	<b>Serafol® AB0</b>	for 10 double determinations or 20 single determinations 1 kit contains 1 twin card, 2 cover foils and 2 mixing sticks.	10 x 1 kit
BG1722		for 50 double determination or 100 single determination clone: A003, B005	50 cards
BG1723	<b>Serafol® AB0+D</b>	for 50 double determinations or 100 single determinations clone: A003, B005, BS226	50 cards
BG1712	<b>Sticks</b>	used to stir each bedside card's reaction field until the reagents are completely dissolved	100 sticks
BG1713	<b>Self-adhesive foil cover</b>	for documentation, the bedside card can be covered by a self-adhesive foil cover	100 foils

## Antigen determination of the ABO system

Art. No.	Product	Description	Packing
BG1101	<b>Anti-A</b>	clone: sifin A-11H5	10 ml
BG1111	<b>Anti-B</b>	clone: sifin B-6F9	10 ml
BG1131	<b>Anti-AB</b>	clone: sifin A-5E10, sifin B-2D7	10 ml
BG1202	<b>Anti-A<sub>1</sub>, lectin</b>	Extract from seeds of <i>Dolichos biflorus</i>	5 ml
BG1212	<b>Anti-H, lectin</b>	Extract from seeds of <i>Laburnum alpinum</i>	5 ml

## Antigen determination of the Rh system

Art. No.	Product	Description	Packing
BG1315	<b>Anti-D (IgM)</b>	clone: BS225	10 ml
BG1354	<b>Anti-C</b>	clone: MS-24, P3X25513G8	5 ml
BG1364	<b>Anti-c</b>	clone: MS-33	5 ml
BG1334	<b>Anti-E</b>	clone: MS-258, 906	5 ml
BG1344	<b>Anti-e</b>	clone: MS-16, MS-21, MS-63	5 ml
BG1371	<b>Anti-C<sup>w</sup></b>	clone: MS-110	2 ml



## Detection of additional blood grouping antigens

Art. No.	Product	Description	Packing
BG1412	<b>Anti-K (Kell)</b>	clone: MS-56	5 ml
BG1421	<b>Anti-k (cellano)</b>	for the anti human globulin test	2 ml
BG3005	<b>Anti-M</b>	clone: LM110, 140	2 ml
BG3006	<b>Anti-N</b>	clone: 20H12, MN879	2 ml

## Anti-Human Globulin (Coombs)

Art. No.	Product	Description	Packing
BG1510	<b>Anti-Human Globulin, polyspecific, dyed</b>	Globulin is used to detect in vivo sensitization of red blood cells in the direct antiglobulin test or to determine blood group antigens and antibodies, respectively (antibody detection test, cross matching) in the indirect antiglobulin test.	10 ml

## Enzymes

Art. No.	Product	Description	Packing
*)	<b>Papain-Cystein</b> *) At the time of going to print, article number and packing were not certain.	Enzyme solution is used for the detection of incomplete antibodies to blood group antigens or for the determination of blood group antigens by the one-stage enzyme test. It may be used in the compatibility test.	*)
BG1613	<b>Bromelin</b>	Enzyme solution is used for the detection of incomplete antibodies to blood group antigens or for the determination of blood group antigens by the one-stage enzyme test. It causes increased agglutination especially in the Rhesus, Kidd, Lewis, Vel, P and ABO systems. It may be used in the compatibility test (cross matching) and in the antibody detection test. Bromelin is also useful in detection of very weak alloagglutinins when determining the ABO blood group.	10 ml

# Blood-Group Determination on the automated Analyser PK7200<sup>®</sup> and PK7300<sup>®</sup>

## Kits for the automated analyser PK7200<sup>®</sup> and PK7300<sup>®</sup>

Art. No.	Product	Description	Packing
BG2000	<b>Kit ABO+D</b>	for 2000 tests Test Reagent Anti-A, clone: A003 Test Reagent Anti-B, clone: B005 Test Reagent Anti-AB, clone: BS63, BS85 Test Reagent Anti-D(1), clone: BS226 Test Reagent Anti-D(2), clone: BS232 Negative control	6 x 50 ml
BG2100	<b>Kit 1 Rh+K</b>	for 2000 tests Test Reagent Anti-C(1), clone: MS-24 Test Reagent Anti-c(1), clone: MS-33 Test Reagent Anti-E(1), clone: MS-258, 906 Test Reagent Anti-e(1), clone: MS-16, MS-21, MS-63 Test Reagent Anti-K(1), clone: MS-56 Negative control	6 x 50 ml
BG2200	<b>Kit 2 Rh+K</b>	for 2000 tests Test Reagent Anti-C(2), clone: MS-273 Test Reagent Anti-c(2), clone: MS-35 Test Reagent Anti-E(2), clone: MS-260, MS-12 Test Reagent-e(2), clone: MS-62, MS-69 Test Reagent Anti-K(2), clone: 601 Negative control	6 x 50 ml

## Test reagents of Kit AB0+D for the automated analyser PK7200® and PK7300®

Art. No.	Product	Description	Packing
BG2010	<b>Test Reagent Anti-A</b>	clone: A003	6 x 50 ml
BG2020	<b>Test Reagent Anti-B</b>	clone: B005	6 x 50 ml
BG2030	<b>Test Reagent Anti-AB</b>	clone: BS63, BS85	6 x 50 ml
BG2040	<b>Test Reagent Anti-D(1)</b>	clone: BS226	6 x 50 ml
BG2050	<b>Test Reagent Anti-D(2)</b>	clone: BS232	6 x 50 ml

## Test reagents of Kit 1 Rh+K for the automated analyser PK7200® and PK7300®

Art. No.	Product	Description	Packing
BG2110	<b>Test Reagent Anti-C(1)</b>	clone: MS-24	6 x 50 ml
BG2120	<b>Test Reagent Anti-c(1)</b>	clone: MS-33	6 x 50 ml
BG2130	<b>Test Reagent Anti-E(1)</b>	clone: MS-258, 906	6 x 50 ml
BG2140	<b>Test Reagent Anti-e(1)</b>	clone: MS-16, MS-21, MS-63	6 x 50 ml
BG2150	<b>Test Reagent Anti-K(1)</b>	clone: MS-56	6 x 50 ml

## Test reagents of Kit 2 Rh+K for the automated analyser PK7200® and PK7300®

Art. No.	Product	Description	Packing
BG2210	<b>Test Reagent Anti-C(2)</b>	clone: MS-273	6 x 50 ml
BG2220	<b>Test Reagent Anti-c(2)</b>	clone: MS-35	6 x 50 ml
BG2230	<b>Test Reagent Anti-E(2)</b>	clone: MS-260, MS-12	6 x 50 ml
BG2240	<b>Test Reagent Anti-e(2)</b>	clone: MS-62, MS-69	6 x 50 ml
BG2250	<b>Test Reagent Anti-K(2)</b>	clone: 601	6 x 50 ml

## Further test reagents for the automated analyser PK7200® and PK7300®

Art. No.	Product	Description	Content
BG2510	<b>Test Reagent Anti-A(2)</b>	clone: sifin A-11H5	6 x 50 ml
BG2520	<b>Test Reagent Anti-B(2)</b>	clone: sifin B-6F9	6 x 50 ml
BG2410	<b>Negative control</b>		6 x 50 ml

# Monoclonal Antibodies for Blood Grouping

## Monoclonal antibodies available as concentrates and bulk

MAB	Clone	Properties	Isotype
Anti-A	<b>A-11H5</b>	Murine monoclonal antibody strongly reacts with red blood cells of the blood groups A <sub>1</sub> , A <sub>2</sub> , A <sub>1</sub> B and A <sub>2</sub> B. Agglutinates A <sub>3</sub> red blood cells (mixed field agglutination), reaction with A <sub>x</sub> red blood cells usually positive.	IgM
Anti-A	<b>A-5E10</b>	Murine monoclonal antibody strongly reacts with red blood cells of the blood groups A <sub>1</sub> , A <sub>2</sub> , A <sub>1</sub> B and A <sub>2</sub> B. Agglutinates A <sub>3</sub> red blood cells (mixed field agglutination), reaction with A <sub>x</sub> red blood cells usually positive.	IgM
Anti-B	<b>B-6F9</b>	Murine monoclonal antibody strongly reacts with red blood cells of the blood groups B, A <sub>1</sub> B and A <sub>2</sub> B. Does not react with „acquired“ B.	IgM
Anti-B	<b>B-2D7</b>	Murine monoclonal antibody strongly reacts with red blood cells of the blood groups B, A <sub>1</sub> B and A <sub>2</sub> B. Does not react with „acquired“ B.	IgM
Anti-D	<b>BS225</b>	Human monoclonal antibody strongly reacts with red blood cells with normally expressed D antigen. Weak D red blood cells are agglutinated subject to the antigen density. Category VI is not detected among the partial D antigens.	IgM
Anti-D	<b>NaTH119</b>	Human monoclonal antibody strongly reacts with red blood cells with normally expressed D antigen. Weak D red blood cells (types 1-4) are agglutinated subject to the antigen density. Categories IV and VI are not detected among the partial D antigens.	IgM
Anti-D	<b>LOR-15C9</b>	Human monoclonal antibody strongly reacts with red blood cells with normally expressed D antigen. Weak D red blood cells (types 1-3) are agglutinated subject to the antigen density. Categories II, III, Va, VI and VII are agglutinated among the partial D antigens.	IgG
Anti-H	<b>A-46/B/B-10</b>	Murine monoclonal antibody strongly reacts with red blood cells of the blood groups O, A <sub>2</sub> and A <sub>2</sub> B positive; with A <sub>1</sub> and A <sub>1</sub> B weak positive or negative.	IgM
Anti-M	<b>M-11H2</b>	Murine monoclonal antibody specifically agglutinates red blood cells with M antigen.	IgG <sub>1</sub>
Anti-N	<b>N-20H12</b>	Murine monoclonal antibody specifically agglutinates red blood cells with N antigen after correction of pH to 8.6 - 8.9.	IgG <sub>2b</sub>

# Bacteriological Test Reagents

DIAGNOSTICS WITH PASSION



**Are you looking for the serological confirmation of suspicious colonies?**

sifin diagnostics gmbh offers a comprehensive line of specific test reagents for serotyping in clinical and veterinary diagnostics. Our tests are based on monoclonal antibodies. This leads to a strong and specific

agglutination without cross reaction.

All our antibodies have been developed in-house and are manufactured in our company. Our products are offered as ready-to-use reagents in dropper bottles or as bulk to IVD companies.

**Our portfolio comprises:**

- Salmonella Diagnostics
- Shigella Diagnostics
- Yersinia Diagnostics
- Coli Diagnostics



# Salmonella Diagnostics

## Test reagents for screening

For the slide agglutination test using polyvalent, group and monospecific test reagents, start with a subculture of the suspect colony or colonies, preferably on non-selective culture medium such as Nutrient Agar, Blood Agar or Kligler Agar.

Before starting the serotyping, it is necessary to confirm biochemically that the isolate belongs to the *Salmonella* genus. For this purpose, alternative procedures (e.g. MALDI mass spectroscopy: MALDI-TOF) can also be used, presuming that the suitability of the procedure has been verified.

Art. No.	Product	Description	Liquid   Lyo.	Packing
TR1101	<b>Anti-Salmonella (A - 67 + Vi), omnivalent</b>	Used for initial testing of suspicious colonies to detect the presence of bacteria of the <i>Salmonella</i> genus.	liquid	1 ml
TR1101-01			liquid	5 ml
TR1111	<b>Anti-Salmonella I (A - E + Vi)</b>	Agglutinates salmonellae of the groups O:2 (A) to O:1,3,19 (E <sub>4</sub> ) and Vi.	liquid	1 ml
TR1111-01			liquid	5 ml
TR1121	<b>Anti-Salmonella II (F - 67)</b>	Agglutinates salmonellae of the groups O:11 (F) to O:67.	lyophilised	1 ml
TR1121-01			lyophilised	5 ml
TR1141	<b>Anti-Salmonella Poly-H Phase 1 &amp; 2</b>	Contains antibodies covering H-antigens or complexes: a, b, c, d, E, G, i, k, L, r, y, Z, Z <sub>4</sub> , Z <sub>6</sub> , Z <sub>10</sub> , Z <sub>29</sub> , Z <sub>35</sub> , Z <sub>38</sub> , Z <sub>41</sub> , H:1.	liquid	1 ml
TR1141-01			liquid	5 ml

## O-Group Pool test reagents

First carry out the test with Anti-Salmonella OMA and Anti-Salmonella OMB: Approx. 98 % of *Salmonella* can be assigned. If the strain does not agglutinate, it is recommended to test this strain with Anti-Salmonella Vi. If this reaction is also negative the strain must be agglutinated with Anti-Salmonella OMC, Anti-Salmonella OMD, Anti-Salmonella OME, Anti-Salmonella OMF and Anti-Salmonella OMG.

Art. No.	Product	Description	Liquid   Lyo.	Packing
TR1151	<b>Anti-Salmonella OMA</b>	(A, B, D, E, L)	liquid	1 ml
TR1151-01			liquid	3 ml
TR1161	<b>Anti-Salmonella OMB</b>	(C, F, G, H)	liquid	1 ml
TR1161-01			liquid	3 ml
TR1170	<b>Anti-Salmonella OMC</b>	(I, J, K, M, N, O, P)	liquid	1 ml
TR1171	<b>Anti-Salmonella OMD</b>	(Q, R, S, T, U, V, W)	liquid	1 ml
TR1172	<b>Anti-Salmonella OME</b>	(X, Y, Z, 51, 52, 53)	liquid	1 ml
TR1173	<b>Anti-Salmonella OMF</b>	(54, 55, 56, 57, 58, 59)	liquid	1 ml
TR1174	<b>Anti-Salmonella OMG</b>	(60, 61, 62, 63, 65, 66, 67)	liquid	1 ml
TR1316	<b>Anti-Salmonella Vi</b>		liquid	1 ml



## H-Phase Pool test reagents

The test, using Anti-Salmonella HMA, Anti-Salmonella HMB or Anti-Salmonella HMC, enables the identification of the most common H-antigens or H-antigen complexes of *Salmonella* strains. Furthermore the presence of the antigen complex H:1 should be tested by Anti-Salmonella H:1 (TR 1437, TR 5437).

Art. No.	Product	Description	Liquid   Lyo.	Packing
TR1181	<b>Anti-Salmonella HMA</b>	(a, b, c, d, i, z <sub>10</sub> , z <sub>29</sub> )	liquid	1 ml
TR1181-01			liquid	3 ml
TR1183	<b>Anti-Salmonella HMB</b>	(E, G)	liquid	1 ml
TR1183-01			liquid	3 ml
TR1185	<b>Anti-Salmonella HMC</b>	(k, y, z, L, Z <sub>4</sub> , r)	liquid	1 ml
TR1185-01			liquid	3 ml

## Group specific test reagents Anti-Salmonella

Mixtures of monoclonal antibodies of the corresponding specificities in the form of cell culture supernatants. To determine the serogroup of *Salmonella* spp. that agglutinate with the omnivalent test reagent and/or the polyspecific test reagent Anti-Salmonella I.

They agglutinate exclusively *Salmonella* spp. of the declared serogroup. According to the frequency of the reported serovars, the group-specific test reagents should be used in the following order:

1. Anti-Salmonella group D
2. Anti-Salmonella group B
3. Anti-Salmonella group C
4. Anti-Salmonella group E
5. Anti-Salmonella O:2 corresponds to Anti-Salmonella group A

Art. No.	Product	Description	Liquid   Lyo.	Packing
TR1201	<b>Anti-Salmonella Group B</b>	Records all antigen combinations of group O:4 (B).	liquid	1 ml
TR1201-01			liquid	5 ml
TR1202	<b>Anti-Salmonella Group C</b>	Records all strains of group O:7 (C <sub>1</sub> ) and O:8 (C <sub>2</sub> -C <sub>3</sub> ).	liquid	1 ml
TR1203	<b>Anti-Salmonella Group D</b>	Records all strains of group O:9 (D <sub>1</sub> ), O:9,46 (D <sub>2</sub> ) and O:9,46,27 (D <sub>3</sub> ).	liquid	1 ml
TR1203-01			liquid	5 ml
TR1204	<b>Anti-Salmonella Group E (O:3 complex)</b>	Records all strains of group O:3,10; O:3,15; O:3,15,34 (E <sub>1</sub> ) and O:1,3,19 (E <sub>2</sub> ).	liquid	1 ml

# Salmonella Diagnostics

## Monospecific test reagents Anti-Salmonella O

For use in identifying specific O-antigens or Vi-antigen. The test reagents for specificities that are needed less often are lyophilised (freeze-dried).

Art. No.	Product	Description	Liquid   Lyo.	Packing
TR1301	<b>Anti-Salmonella O:2</b>	group A	liquid	1 ml
TR1302	<b>Anti-Salmonella O:4</b>	group B	liquid	1 ml
TR1302-01			liquid	5 ml
TR1303	<b>Anti-Salmonella O:5</b>	group B	liquid	1 ml
TR1303-01			liquid	5 ml
TR1304	<b>Anti-Salmonella O:6<sub>1</sub></b>	group C <sub>2</sub> -C <sub>3</sub>	lyophilised	1 ml
TR1305	<b>Anti-Salmonella O:7</b>	group C <sub>1</sub>	liquid	1 ml
TR1306	<b>Anti-Salmonella O:8</b>	group C <sub>2</sub> -C <sub>3</sub>	liquid	1 ml
TR1307	<b>Anti-Salmonella O:9</b>	group D	liquid	1 ml
TR1307-01			liquid	5 ml
TR1308	<b>Anti-Salmonella O:10</b>	group E <sub>1</sub> and group E <sub>4</sub>	liquid	1 ml
TR1323	<b>Anti-Salmonella O:11</b>	group F	lyophilised	1 ml
TR1325	<b>Anti-Salmonella O:13</b>	group G	lyophilised	1 ml
TR1309	<b>Anti-Salmonella O:14</b>	group H, not group C <sub>1</sub>	lyophilised	1 ml
TR1310	<b>Anti-Salmonella O:15</b>	group E <sub>1</sub>	liquid	1 ml
TR1328	<b>Anti-Salmonella O:16</b>	group I	lyophilised	1 ml
TR1329	<b>Anti-Salmonella O:17</b>	group J	lyophilised	1 ml
TS1330	<b>Anti-Salmonella O:18</b>	group K	lyophilised	1 ml
TR1311	<b>Anti-Salmonella O:19</b>	group E <sub>4</sub>	liquid	1 ml
TR1312	<b>Anti-Salmonella O:20</b>	group C <sub>2</sub> -C <sub>3</sub>	liquid	1 ml
TR1331	<b>Anti-Salmonella O:21</b>	group L	lyophilised	1 ml
TS1332	<b>Anti-Salmonella O:22</b>	group G	lyophilised	1 ml
TR1335	<b>Anti-Salmonella O:25</b>	group H	lyophilised	1 ml
TR1313	<b>Anti-Salmonella O:27</b>	group B and group D <sub>3</sub>	liquid	1 ml
TR1336	<b>Anti-Salmonella O:28</b>	group M	lyophilised	1 ml
TR1339	<b>Anti-Salmonella O:30</b>	group N	lyophilised	1 ml
TR1314	<b>Anti-Salmonella O:34</b>	group E <sub>3</sub>	liquid	1 ml
TR1341	<b>Anti-Salmonella O:35</b>	group O	lyophilised	1 ml
TR1344	<b>Anti-Salmonella O:38</b>	group P	lyophilised	1 ml
TR1345	<b>Anti-Salmonella O:39</b>	group Q	lyophilised	1 ml

Art. No.	Product	Description	Liquid   Lyo.	Packing
TR1346	<b>Anti-Salmonella O:40</b>	group R	lyophilised	1 ml
TR1347	<b>Anti-Salmonella O:41</b>	group S	lyophilised	1 ml
TR1348	<b>Anti-Salmonella O:42</b>	group T	lyophilised	1 ml
TR1349	<b>Anti-Salmonella O:43</b>	group U	lyophilised	1 ml
TR1350	<b>Anti-Salmonella O:44</b>	group V	lyophilised	1 ml
TR1351	<b>Anti-Salmonella O:45</b>	group W	lyophilised	1 ml
TR1315	<b>Anti-Salmonella O:46</b>	group D <sub>2</sub>	liquid	1 ml
TR1353	<b>Anti-Salmonella O:47</b>	group X	lyophilised	1 ml
TR1354	<b>Anti-Salmonella O:48</b>	group Y	lyophilised	1 ml
TR1355	<b>Anti-Salmonella O:50</b>	group Z	lyophilised	1 ml
TR1356	<b>Anti-Salmonella O:51</b>		lyophilised	1 ml
TR1357	<b>Anti-Salmonella O:52</b>		lyophilised	1 ml
TR1358	<b>Anti-Salmonella O:53</b>		lyophilised	1 ml
TR1359	<b>Anti-Salmonella O:54</b>		lyophilised	1 ml
TR1360	<b>Anti-Salmonella O:55</b>		lyophilised	1 ml
TR1361	<b>Anti-Salmonella O:56</b>		lyophilised	1 ml
TR1362	<b>Anti-Salmonella O:57</b>		lyophilised	1 ml
TR1363	<b>Anti-Salmonella O:58</b>		lyophilised	1 ml
TR1364	<b>Anti-Salmonella O:59</b>		lyophilised	1 ml
TR1365	<b>Anti-Salmonella O:60</b>		lyophilised	1 ml
TR1366	<b>Anti-Salmonella O:61</b>		lyophilised	1 ml
TR1367	<b>Anti-Salmonella O:62</b>		lyophilised	1 ml
TR1368	<b>Anti-Salmonella O:63</b>		lyophilised	1 ml
TR1369	<b>Anti-Salmonella O:65</b>		lyophilised	1 ml
TR1370	<b>Anti-Salmonella O:66</b>		lyophilised	1 ml
TR1371	<b>Anti-Salmonella O:67</b>		lyophilised	1 ml

## Monospecific test reagent Anti-Salmonella Vi

Identification of the Vi-antigen.

Art. No.	Product	Liquid   Lyo.	Packing
TR1316	<b>Anti-Salmonella Vi</b>	liquid	1 ml

# Salmonella Diagnostics

## Monospecific test reagents Anti-Salmonella H

For use in identifying or checking the H-antigens or H-antigen complexes of *Salmonella* strains in accordance with the White-Kauffmann-Le-Minor scheme using slide agglutination. They enable the serovar to be determined.

Art. No.	Product	Phase induction	Liquid   Lyo.	Packing
TR1401	<b>Anti-Salmonella H:a</b>	x	liquid	1 ml
TR1402	<b>Anti-Salmonella H:b</b>	x	liquid	1 ml
TR1403	<b>Anti-Salmonella H:c</b>	x	liquid	1 ml
TR1404	<b>Anti-Salmonella H:d</b>	x	liquid	1 ml
TR1405	<b>Anti-Salmonella H:E</b>	x	liquid	1 ml
TR1405-01		x	liquid	5 ml
TR1407	<b>Anti-Salmonella H:f</b>		lyophilised	1 ml
TR1406	<b>Anti-Salmonella H:g</b>	x	liquid	1 ml
TR1406-01		x	liquid	5 ml
TR1408	<b>Anti-Salmonella H:g,m</b>	x	liquid	1 ml
TR1408-01		x	liquid	5 ml
TR1409	<b>Anti-Salmonella H:h</b>		liquid	1 ml
TR1410	<b>Anti-Salmonella H:i</b>	x	liquid	1 ml
TR1410-01		x	liquid	5 ml
TR1411	<b>Anti-Salmonella H:k</b>	x	lyophilised	1 ml
TR1412	<b>Anti-Salmonella H:l</b>	x	liquid	1 ml
TR1412-01		x	liquid	5 ml
TS1413	<b>Anti-Salmonella H:m</b>		lyophilised	1 ml
TR1438	<b>Anti-Salmonella H:n</b>	x	liquid	1 ml
TS1414	<b>Anti-Salmonella H:p</b>		lyophilised	1 ml
TS1415	<b>Anti-Salmonella H:q</b>		lyophilised	1 ml
TR1416	<b>Anti-Salmonella H:r</b>	x	liquid	1 ml
TS1417	<b>Anti-Salmonella H:s</b>		lyophilised	1 ml
TS1418	<b>Anti-Salmonella H:t</b>		lyophilised	1 ml
TS1419	<b>Anti-Salmonella H:u</b>		lyophilised	1 ml
TS1420	<b>Anti-Salmonella H:v</b>		lyophilised	1 ml
TS1421	<b>Anti-Salmonella H:w</b>		lyophilised	1 ml
TS1422	<b>Anti-Salmonella H:x</b>		lyophilised	1 ml
TR1423	<b>Anti-Salmonella H:y</b>	x	liquid	1 ml
TR1424	<b>Anti-Salmonella H:z</b>	x	liquid	1 ml
TS1425	<b>Anti-Salmonella H:z<sub>4r</sub>z<sub>23</sub></b>		lyophilised	1 ml

Art. No.	Product	Description	Phase induction	Liquid   Lyo.	Packing
TS1426	<b>Anti-Salmonella H:z<sub>6</sub></b>			lyophilised	1 ml
TR1427	<b>Anti-Salmonella H:z<sub>10</sub></b>		x	liquid	1 ml
TR1439	<b>Anti-Salmonella H:z<sub>13</sub></b>	for use by reference labs		liquid	1 ml
TS1428	<b>Anti-Salmonella H:z<sub>15</sub></b>			lyophilised	1 ml
TR1440	<b>Anti-Salmonella H:z<sub>23</sub></b>			lyophilised	1 ml
TS1429	<b>Anti-Salmonella H:z<sub>24</sub></b>			lyophilised	1 ml
TS1449	<b>Anti-Salmonella H:z<sub>28</sub></b>			lyophilised	1 ml
TS1430	<b>Anti-Salmonella H:z<sub>29</sub></b>			lyophilised	1 ml
TS1431	<b>Anti-Salmonella H:z<sub>32</sub></b>			lyophilised	1 ml
TR1445	<b>Anti-Salmonella H:z<sub>35</sub></b>		x	lyophilised	1 ml
TR1447	<b>Anti-Salmonella H:z<sub>38</sub></b>		x	lyophilised	1 ml
TR1448	<b>Anti-Salmonella H:z<sub>41</sub></b>		x	lyophilised	1 ml
TR1437	<b>Anti-Salmonella H:1</b>		x	liquid	1 ml
TR1437-01			x	liquid	5 ml
TR1433	<b>Anti-Salmonella H:2</b>			lyophilised	1 ml
TR1433-01				lyophilised	5 ml
TS1434	<b>Anti-Salmonella H:5</b>			lyophilised	1 ml
TR1435	<b>Anti-Salmonella H:6</b>			lyophilised	1 ml
TS1436	<b>Anti-Salmonella H:7</b>			lyophilised	1 ml

## Control antigens for the Anti-Salmonella test reagents

The control antigens are used to check the agglutinability of the Anti-Salmonella test reagents and for quality control when carrying out the slide agglutination test.

Art. No.	Product	Antigen formula	Liquid   Lyo.	Packing
TS1501	<b>Control antigen Salmonella Paratyphi A-OH</b>	2,12:a:[1,5] (Usually occurs as a monophasic variant.)	liquid	5 ml
TS1502	<b>Control antigen Salmonella Paratyphi B-OH</b>	1,4,[5],12:b:1,2	liquid	5 ml
TS1503	<b>Control antigen Salmonella Paratyphi C-OH</b>	6,7,[Vi]:c:1,5 (Use TS 1507 for Vi.)	liquid	5 ml
TS1504	<b>Control antigen Salmonella Typhi-OH</b>	9,12, [Vi]:d- (Use TS 1507 for Vi.)	liquid	5 ml
TS1505	<b>Control antigen Salmonella Typhimurium-OH</b>	1,4,[5],12:i:1,2 (Often occurs as a monophasic variant.)	liquid	5 ml
TS1506	<b>Control antigen Salmonella Enteritidis-OH</b>	1,9,12:g,m:-	liquid	5 ml
TS1507	<b>Control antigen Salmonella Vi</b>		liquid	5 ml

# Salmonella Diagnostics

## Salmonella O test antigens for the Widal reaction

For use in establishing the existence and determining the quantity of specific agglutinating *Salmonella* antibodies (or agglutinins) in human sera or sera of other origin in the Widal reaction. The test can be performed in test tubes or on a microtitre plate. The test antigens must be diluted 1:10 in PBS.

Art. No.	Product	Description	Liquid   Lyo.	Packing
TS1606	<b>Paratyphi A-O test antigen</b>	(2,12)	liquid	10 ml
TS1601	<b>Paratyphi B-O test antigen</b>	(1,4,5,12)	liquid	10 ml
TS1607	<b>Paratyphi C-O test antigen</b>	(6,7)	liquid	10 ml
TS1602	<b>Typhi O test antigen</b>	(9,12)	liquid	10 ml

## Salmonella OH test antigens for the Widal reaction

The test with OH-test antigens can be performed in test tubes or on a microtitre plate. The test antigens must be diluted 1:10 in PBS.

Art. No.	Product	Description	Liquid   Lyo.	Packing
TS1613	<b>Paratyphi A-OH test antigen</b>	(2,12:a:-)	liquid	10 ml
TS1614	<b>Paratyphi B-OH test antigen</b>	(1,4,5,12:b:1,2)	liquid	10 ml
TS1615	<b>Paratyphi C-OH test antigen</b>	(6,7:c:1,5)	liquid	10 ml
TS1616	<b>Typhi OH test antigen</b>	(9,12:d:-)	liquid	10 ml
TS1611	<b>Typhimurium OH test antigen</b>	(1,4,5,12:i:1,2)	liquid	10 ml
TS1612	<b>Enteritidis OH test antigen</b>	(1,9,12:g,m:-)	liquid	10 ml

## Salmonella H test antigens for the Widal reaction

H-test antigens are suitable only for the test tube test. The test antigens must be diluted 1:10 in PBS.

Art. No.	Product	Liquid   Lyo.	Packing
TS1631	<b>H:a test antigen</b>	liquid	10 ml
TS1632	<b>H:b test antigen</b>	liquid	10 ml
TS1633	<b>H:c test antigen</b>	liquid	10 ml
TS1603	<b>H:d test antigen</b>	liquid	10 ml



## Control sera for the *Salmonella* O and OH test antigens

For use in system control and in checking the agglutinability of the *Salmonella* test antigens in the Widal reaction. The test can be performed in test tubes or on a microtitre plate.

Art. No.	Product	Packing
TS1626	<b>Anti-Salmonella Paratyphi A</b>	1 ml
TS1604	<b>Anti-Salmonella Paratyphi B</b>	1 ml
TS1627	<b>Anti-Salmonella Paratyphi C</b>	1 ml
TS1605	<b>Anti-Salmonella Typhi</b>	1 ml
TS1624	<b>Anti-Salmonella Typhimurium</b>	1 ml
TS1625	<b>Anti-Salmonella Enteritidis</b>	1 ml

## Control sera for the *Salmonella* H test antigens

For use in system control and in checking the agglutinability of the *Salmonella* test antigens in the Widal reaction. The test can be performed in test tubes only.

Art. No.	Product	Description	Packing
TS1641	<b>Anti-Salmonella H:a</b>	(2,12)	1 ml
TS1642	<b>Anti-Salmonella H:b</b>	(1,4,5,12)	1 ml
TS1643	<b>Anti-Salmonella H:c</b>	(6,7)	1 ml
TS1644	<b>Anti-Salmonella H:d</b>	(9,12)	1 ml

# Shigella Diagnostics

## Polyspecific test reagents Anti-Shigella

Serological detection of the serovar of *Shigella* strains using the slide agglutination test.

Art. No.	Product	Contains antibodies against	Liquid   Lyo.	Packing
TR1811	<b>Anti-Shigella I</b>	<i>S. flexneri</i> , type 1 to 6, group 3,4 (y), 6 and 7,8 (x) and <i>S. sonnei</i> S and F form (phase I and II)	lyophilised	1 ml
TR1811-01			lyophilised	5 ml
TS1821	<b>Anti-Shigella II</b>	<i>S. dysenteriae</i> , type 1 to 10 (subgroup A)	lyophilised	1 ml
TS1821-01			lyophilised	5 ml
TS1831	<b>Anti-Shigella III</b>	<i>S. boydii</i> , type 1 to 15 (subgroup C)	lyophilised	1 ml
TS1901	<b>Anti-Shigella flexneri</b>	type 1 - 6 and group 3,4 (y), 6 and 7,8 (x) (subgroup B)	lyophilised	1 ml

## Monospecific test reagents Anti-Shigella

Serological detection and the determination of the serovar of *Shigella* O antigens using the slide agglutination test.

Art. No.	Product	Description	Liquid   Lyo.	Packing
TS2001	<b>Anti-Shigella dysenteriae type 1</b>	monospecific	lyophilised	1 ml
TS2002	<b>Anti-Shigella dysenteriae type 2</b>	monospecific	lyophilised	1 ml
TS2003	<b>Anti-Shigella flexneri type 1</b>	monospecific	lyophilised	1 ml
TS2004	<b>Anti-Shigella flexneri type 2</b>	monospecific	lyophilised	1 ml
TS2005	<b>Anti-Shigella flexneri type 3</b>	monospecific	lyophilised	1 ml
TS2006	<b>Anti-Shigella flexneri type 4</b>	monospecific	lyophilised	1 ml
TS2007	<b>Anti-Shigella flexneri type 5</b>	monospecific	lyophilised	1 ml
TS2008	<b>Anti-Shigella flexneri type 6</b>	monospecific	lyophilised	1 ml
TS2009	<b>Anti-Shigella flexneri group 3,4 (y)</b>	monospecific	liquid	1 ml
TS2010	<b>Anti-Shigella flexneri group 6</b>	monospecific	lyophilised	1 ml
TS2011	<b>Anti-Shigella flexneri group 7,8 (x)</b>	monospecific	lyophilised	1 ml
TR2012	<b>Anti-Shigella sonnei S form</b>	phase I, monospecific	lyophilised	1 ml
TR2013	<b>Anti-Shigella sonnei F form</b>	phase II, monospecific	lyophilised	1 ml
TR2014	<b>Anti-Shigella sonnei S form and F form</b>	phase I and phase II, monospecific (subgroup D)	lyophilised	1 ml

## Control antigens for the Anti-Shigella test reagents

The control antigens are used to check the agglutinability of the Anti-Shigella test reagents and for quality control when carrying out the slide agglutination test.

Art. No.	Product	Description	Liquid   Lyo.	Packing
TS1510	<b>Control antigen Shigella flexneri</b>	<i>S. flexneri</i> Type 1b, 2a, 3a, 4a	liquid	2 ml
TS1511	<b>Control antigen Shigella dysenteriae</b>	<i>S. dysenteriae</i> Type 1, 2, 3, 7	liquid	2 ml
TS1512	<b>Control antigen Shigella boydii</b>	<i>S. boydii</i> Type 1, 2, 5, 8	liquid	2 ml
TS1513	<b>Control antigen Shigella sonnei</b>	<i>S. sonnei</i> S-form (phase I) <i>S. sonnei</i> F-form (phase II)	liquid	2 ml

# Yersinia Diagnostics

## Monospecific test reagents Anti-Yersinia enterocolitica O

Identification of human pathogenic serovars of the species *Yersinia enterocolitica* by slide agglutination.

Art. No.	Product	Liquid   Lyo.	Packing
TS1701	<b>Anti-Yersinia enterocolitica O 3</b>	lyophilised	1 ml
TS1704	<b>Anti-Yersinia enterocolitica O 5</b>	lyophilised	1 ml
TS1705	<b>Anti-Yersinia enterocolitica O 8</b>	lyophilised	1 ml
TS1703	<b>Anti-Yersinia enterocolitica O 9</b>	lyophilised	1 ml
TS1706	<b>Anti-Yersinia enterocolitica O 27</b>	lyophilised	1 ml

SALE

## Yersinia enterocolitica O test antigens for the Widal reaction

For use in detecting the existence and determining the quantity of specific agglutinating *Yersinia enterocolitica* O antibodies (or agglutinins) in human sera or sera of other origin using the Widal reaction. The test is performed on a microtitre plate with dyed antigen.

Art. No.	Product	Liquid   Lyo.	Packing
TS1721	<b>Yersinia enterocolitica-O test antigen (3)</b>	lyophilised	3 ml
TS1724	<b>Yersinia enterocolitica-O test antigen (5)</b>	lyophilised	3 ml
TS1723	<b>Yersinia enterocolitica-O test antigen (9)</b>	lyophilised	3 ml

## Control sera Anti-Yersinia for the Widal reaction

For use in system control and in checking the agglutinability of the *Yersinia enterocolitica* test antigens in the Widal reaction. The test is performed on a microtitre plate.

Art. No.	Product	Liquid   Lyo.	Packing
TS1733	<b>Anti-Yersinia enterocolitica O 3</b>	lyophilised	1 ml
TS1734	<b>Anti-Yersinia enterocolitica O 5</b>	lyophilised	1 ml
TS1735	<b>Anti-Yersinia enterocolitica O 9</b>	lyophilised	1 ml

# Coli Diagnostics

## Polyspecific test reagents Anti-Coli

Serological detection of isolated *E. coli* strains from human test material or other origin by slide agglutination.

Art. No.	Product	Contains antibodies against	Liquid   Lyo.	Packing
TS2111	<b>Anti-Coli I</b>	O 26:K 60, O 44:K 74, O 114:K90,	lyophilised	1 ml
TS2111-01		O 125:K 70, O 142:K 86, O 158:K -	lyophilised	5 ml
TR2121	<b>Anti-Coli II</b>	O 55:K 59, O 86:K 61, O 91:K -,	lyophilised	1 ml
TR2121-01		O 111:K 58, O 119:K 69, O 126:K 71, O 127:K 63, O 128:K 67	lyophilised	5 ml
TR2131	<b>Anti-Coli III</b>	O 25:K 11, O 78:K 80, O 103:K -,	lyophilised	1 ml
TR2131-01		O 118:K -, O 124:K 72, O 145:K -, O 157:K -, O 164:K -	lyophilised	5 ml

# Coli Diagnostics

## Monospecific test reagents Anti-Coli

Used for the serological detection and serovar determination of isolated *E. coli* strains from human test material or other origin by slide agglutination and Widal reaction (confirmation test).

Art. No.	Product	Liquid   Lyo.	Packing
TS2201	<b>Anti-Coli O 25:K 11</b>	lyophilised	1 ml
TS2202	<b>Anti-Coli O 26:K 60</b>	lyophilised	1 ml
TS2203	<b>Anti-Coli O 44:K 74</b>	lyophilised	1 ml
TS2204	<b>Anti-Coli O 55:K 59</b>	lyophilised	1 ml
TR2205	<b>Anti-Coli O 78:K 80</b>	lyophilised	1 ml
TS2206	<b>Anti-Coli O 86:K 61</b>	lyophilised	1 ml
TS2222	<b>Anti-Coli O 91:K -</b>	lyophilised	1 ml
TS2216	<b>Anti-Coli O 103:K -</b>	lyophilised	1 ml
TS2207	<b>Anti-Coli O 111:K 58</b>	lyophilised	1 ml
TS2208	<b>Anti-Coli O 114:K 90</b>	lyophilised	1 ml
TS2220	<b>Anti-Coli O 118:K -</b>	lyophilised	1 ml
TS2209	<b>Anti-Coli O 119:K 69</b>	lyophilised	1 ml
TR2210	<b>Anti-Coli O 124:K 72</b>	lyophilised	1 ml
TS2211	<b>Anti-Coli O 125:K 70</b>	lyophilised	1 ml
TS2212	<b>Anti-Coli O 126:K 71</b>	lyophilised	1 ml
TS2213	<b>Anti-Coli O 127:K 63</b>	lyophilised	1 ml
TS2214	<b>Anti-Coli O 128:K 67</b>	lyophilised	1 ml
TS2215	<b>Anti-Coli O 142:K 86</b>	lyophilised	1 ml
TS2221	<b>Anti-Coli O 145:K -</b>	lyophilised	1 ml
TR2218	<b>Anti-Coli O 157:K -</b>	lyophilised	1 ml
TS2219	<b>Anti-Coli O 158:K -</b>	lyophilised	1 ml
TS2217	<b>Anti-Coli O 164:K -</b>	lyophilised	1 ml



# E. coli relevant to veterinary medicine

## Coli diagnostics in young poultry

The polyspecific screening reagent Anti-Coli A is intended for use in the serological detection of O 1, O 2, O 18 as well as O 78 antigen of *E. coli* strains isolated from test material, using slide agglutination. An isolate with positive results requires testing with monospecific reagents for further characterisation and to rule out unspecific agglutinations a check by the Gruber-Widal test (confirmation test).

Art. No.	Product	Description	Liquid   Lyo.	Packing
TR2311	<b>Anti-Coli A</b>	polyspecific (O 1, O 2, O 18, O 78) serological detection of O 1-, O 2-, O 18 or O 78 antigen.	lyophilised	1 ml
TS2401	<b>Anti-Coli O 1</b>	monospecific	lyophilised	1 ml
TS2501	<b>Gruber Widal Serum Anti-Coli O 1</b>	for the confirmation test	lyophilised	1 ml
TS2402	<b>Anti-Coli O 2</b>	monospecific	lyophilised	1 ml
TS2502	<b>Gruber Widal Serum Anti-Coli O 2</b>	for the confirmation test	lyophilised	1 ml
TS2403	<b>Anti-Coli O 18</b>	monospecific	lyophilised	1 ml
TS2503	<b>Gruber Widal Serum Anti-Coli O 18</b>	for the confirmation test	lyophilised	1 ml
TR2205	<b>Anti-Coli O 78:K 80</b>	monospecific	lyophilised	1 ml

## Coli diagnostics in young cattles

The test sera are used to test for the presence of type-specific antigens with *Escherichia coli* isolates from cattles. They are used as evidence of antigens to the cell surface of the *E. coli* isolate (O antigen, K antigen and F antigen) using slide agglutination. First carry out a screening agglutination with the polyspecific test serum Anti-Coli C. If the reaction with the polyspecific test serum is positive, the isolate must then be typed with the monospecific test sera. The F5 (K99) antigen is often not formed in sufficient quantities on standard culture media. We therefore recommend the use of Minca Agar when culturing to promote the formation of the fimbrial antigen F5 (K99).

Art. No.	Product	Description	Liquid   Lyo.	Packing
TS2601	<b>Anti-Coli C</b>	polyspecific (O 9:K 35, O 101:K 28, O 101:K 30, O 101:K 32) Contains antibodies directed against the <i>E. coli</i> types listed as monospecific (TS 2611, TS 2612, TS 2613, TS 2614) and the fimbrial antigen F 5 (K 99).	liquid	1 ml
TS2615	<b>Anti-Coli F 5 (K 99)</b>	monospecific	liquid	1 ml
TS2611	<b>Anti-Coli O 9:K 35</b>	monospecific	liquid	1 ml
TS2612	<b>Anti-Coli O 101:K 28</b>	monospecific	liquid	1 ml
TS2613	<b>Anti-Coli O 101:K 30</b>	monospecific	liquid	1 ml
TS2614	<b>Anti-Coli O 101:K 32</b>	monospecific	liquid	1 ml
TR2205	<b>Anti-Coli O 78:K 80</b>	monospecific	lyophilised	1 ml
TN1040	<b>Minca Agar, modified</b>	see also dehydrated culture media		500 g
TN1334	<b>Minca Supplement</b>	see also supplements		12 x 1 vial

# E. coli relevant to veterinary medicine

## Coli diagnostics in young pigs

The test sera are used to test for the presence of type-specific antigens with *Escherichia coli* isolates from pigs. They are used as evidence of antigens to the cell surface of the *E. coli* isolate (O antigen, K antigen and F antigen) using slide agglutination (Serotyping). First carry out a screening agglutination with the polyspecific test serum Anti-Coli P. If the reaction with the polyspecific test serum is positive, the isolate must then be typed with the monospecific test sera.

Art. No.	Product	Description	Liquid   Lyo.	Packing
TS2701	<b>Anti-Coli P</b>	polyspecific (O 8:K 87, O 138:K 81, O 139:K 82, O 141:K 85, O 147:K 89, O 149:K 91) Contains antibodies directed against the <i>E. coli</i> types listed as monospecific (TS 2711, TS 2712, TS 2713, TS 2714, TS 2715, TS 2716) and the fimbrial antigen F 4 (K 88).	liquid	1 ml
TS2717	<b>Anti-Coli F 4 (K 88)</b>	monospecific	liquid	1 ml
TS2711	<b>Anti-Coli O 8:K 87</b>	monospecific	liquid	1 ml
TS2712	<b>Anti-Coli O 138:K 81</b>	monospecific	liquid	1 ml
TS2713	<b>Anti-Coli O 139:K 82</b>	monospecific	liquid	1 ml
TS2714	<b>Anti-Coli O 141:K 85</b>	monospecific	liquid	1 ml
TS2715	<b>Anti-Coli O 147:K 89</b>	monospecific	liquid	1 ml
TS2716	<b>Anti-Coli O 149:K 91</b>	monospecific	liquid	1 ml

# Culture Media

DIAGNOSTICS WITH PASSION



## Culture media at first-hand

We are manufacturer of more than 200 different culture media and supplements in order to cultivate the microorganisms or specifically promote the growth of particularly fastidious species.

Thus, you are optimally equipped for the analysis of veterinary or

human samples. Special media for food, water, pharmaceutical products complete the portfolio.

Do you have a specific request? Or are you looking for a supplier of media based on your own formulation?

We develop and produce according to your demand.

## Our portfolio comprises:

- Dehydrated culture media
- Supplements
- Additives
- Detection reagents
- Ready-to-use culture media
- Base materials



# Dehydrated Culture Media

## A

Art. No.	Product	Description	Packing
TN1294	<b>Alkaline Peptone Water with 2 % Sodium Chloride (APW)</b>	for 15.2 l culture medium (33.0 g for 1 l culture medium) Enrichment of pathogenic vibrios.	500 g
TN1103	<b>Azide Glucose Broth</b>	for 7.4 l culture medium: single strength (70.0 g for 1 l medium) Cultivation of enterococci from water (acc. to § 64 German Food and Feed Code) and waste water.	500 g

## B

Art. No.	Product	Description	Packing
TN1245	<b>Bacillus Cereus Agar (Base) (PEMBA)</b>	for 12.5 l culture medium (39.9 g for 1 l culture medium) Isolation and colony count of <i>Bacillus cereus</i> in foodstuffs. Addition of Egg Yolk Emulsion TN 1316 and Bacillus Cereus Selective Supplement TN 1315.	500 g
TN1104	<b>Baird Parker Agar (Base)</b>	for 7.9 l culture medium (63.0 g for 1 l culture medium) Isolation and differentiation of <i>Staphylococcus aureus</i> from foodstuffs (acc. to § 64 German Food and Feed Code). Addition of Egg Yolk Tellurite Emulsion TN 1310.	500 g
TN1133 TN1133-01	<b>Bile Chrysoidine Glycerol Agar (Base) (GCG Agar)</b>	for 11.7 l culture medium (42.8 g for 1 l culture medium) for 116.8 l culture medium (42.8 g for 1 l culture medium) Isolation and differentiation of <i>Enterobacteriaceae</i> and various aerobic gram-negative bacteria from clinical material (especially urinary diagnostics and „Variadiagnostik“). Addition of urea TN 1308 and glycerin TN 1424.	500 g 5 kg
TN1237	<b>Bile Chrysoidine Glycerol Agar (Base) with MUG</b>	for 11.7 l culture medium (42.9 g for 1 l culture medium) Isolation and differentiation of <i>Enterobacteriaceae</i> and various aerobic gram-negative bacteria from clinical material. Advantage: $\beta$ -D-glucuronidase-positive <i>E. coli</i> strains are easy to detect by fluorescence. Addition of urea TN 1308 and Glycerin TN 1424.	500 g
TN1276	<b>Bile Esculin Azide Agar (BEM, Bile Esculin Medium)</b>	for 9.3 l culture medium (53.7 g for 1 l culture medium) Selective detection and determination of the germ count of enterococci from water and other material.	500 g
TN1106	<b>Blood Agar (Base)</b>	for 13.9 l culture medium (36.0 g for 1 l culture medium) Used to produce blood plates and boiled blood plates, for the isolation and breeding of various fastidious and above all pathogenic microorganisms, and for determining their haemolysis forms. Addition of blood.	500 g

Art. No.	Product	Description	Packing
TN1216	<b>Brain Heart Infusion Broth (BHI Broth, Brain Heart Glucose Broth)</b>	for 13.5 l culture medium (37.0 g for 1 l medium) Cultivation of various fastidious microorganisms (acc. to § 64 German Food and Feed Code).	500 g
TN1109	<b>Brilliant Green Bile Lactose Broth (BRILA Broth)</b>	for 12.5 l culture medium: single strength (40.0 g for 1 l medium) for 6.3 l culture medium: double strength (80.0 g for 1 l medium) Selective enrichment of <i>Escherichia coli</i> and other coliforms in water, dairy products, foodstuffs (acc. to § 64 German Food and Feed Code) and from other test materials. Culture medium is used for confirmation of <i>E. coli</i> and coliform bacteria in accordance with IDF-Standard 73B:1998.	500 g
TN1110	<b>Brilliant Green Phenol Red Agar acc. to Edel and Kampelmacher (BPLS Agar, modified)</b>	for 10.1 l culture medium (49.7 g for 1 l culture medium) Selective culture medium for use of <i>Salmonella</i> from foodstuffs (acc. to § 64 German Food and Feed Code).	500 g
TN1111	<b>Brilliant Green Phenol Red Agar (BPLS Agar acc. to Kauffmann, modified)</b>	for 10.0 l culture medium (50.0 g for 1 l culture medium) Isolation and identification of salmonellae (with the exception of <i>S. Typhi</i> ) from meat and other foodstuffs (acc. to § 64 German Food and Feed Code) as well as for the examination of pharmaceutical preparations.	500 g
TN1113	<b>Bromothymol Blue Broth (Base)</b>	for 33.3 l culture medium (15.0 g for 1 l culture medium) Culture medium for testing carbohydrate utilisation after addition of carbohydrates, especially by fast-growing bacteria, e.g. <i>Enterobacteriaceae</i> .	500 g
TN1078	<b>Buffered Nitrate Motility Medium</b>	for 25.0 l culture medium (20.0 g for 1 l culture medium) Confirmation of <i>Clostridium perfringens</i> (acc. to § 64 German Food and Feed Code). Addition of glycerin TN 1424.	500 g
TN1137 TN1137-01	<b>Buffered Peptone Water (BPW)</b>	for 25.0 l culture medium (20.0 g for 1 l culture medium) for 250.0 l culture medium (20.0 g for 1 l culture medium) Pre-enrichment of salmonellae from milk, dairy products and of other microorganisms from other foodstuffs.	500 g 5 kg
TN1025 TN1025-01	<b>Buffered Sodium Chloride Peptone Solution pH 7.0 acc. to harm. EP/USP/JP</b>	for 34.2 l culture medium (14.6 g for 1 l culture medium) for 342.4 l culture medium (14.6 g for 1 l culture medium) Dilution fluid for samples in case of microbiological contamination. Culture medium for testing non-sterile products based on the recommendations of the harmonised method acc. to EP/USP/JP (2006).	500 g 5 kg

# Dehydrated Culture Media

## C

Art. No.	Product	Description	Packing
TN1031 TN1031-01	<b>Casein Soya Bean Digest Agar acc. to harm. EP/USP/JIP (CASO Agar acc. to harm. EP/USP/JIP)</b>	for 12.5 l culture medium (40.0 g for 1 l culture medium) for 125.0 l culture medium (40.0 g for 1 l culture medium) Universal culture medium, inhibitor-free and indicator-free, for a wide range of applications. Culture medium for testing non-sterile products based on the recommendations of the harmonised method acc. to EP/USP/JIP (2006).	500 g 5 kg
TN1263 TN1263-01	<b>Casein Soya Bean Digest Broth acc. to harm. EP/USP/JIP (CASO Broth acc. to harm. EP/USP/JIP)</b>	for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) Universal culture medium for the cultivation of bacteria and fungi. Suitable for cold filtration. Culture medium for testing non-sterile products based on the recommendations of the harmonised method acc. to EP/USP/JIP (2006).	500 g 5 kg
TN1273	<b>CATC Agar (Base) (Citrate Azide Tween Carbonate Agar)</b>	for 9.2 l culture medium (54.5 g for 1 l culture medium) Isolation of enterococci from foodstuffs. Addition of Tween® 80 TN 1422.	500 g
TN1024	<b>Cetrimide Agar acc. to harm. EP/USP/JIP</b>	for 11.0 l culture medium (45.3 g for 1 l culture medium) Test for <i>Pseudomonas aeruginosa</i> . Culture medium for testing non-sterile products. This culture medium is based on the recommendations of the harmonised method acc. to EP/USP/JIP (2006). Addition of glycerin TN 1424.	500 g
TN1114	<b>China-blue Lactose Agar</b>	for 13.0 l culture medium (38.5 g for 1 l culture medium) Determination of the colony count and for the differentiation of bacteria from milk.	500 g
TN1064	<b>ChroMedium Coliform (Chromogenic Coliform Agar, CCA)</b>	for 17.3 l culture medium (28.9 g for 1 l culture medium) Chromogenic selective agar to detect <i>E. coli</i> and coliform bacteria from water with low accompanying flora (acc. to the draft standard DIN EN ISO 9308-1:2012) and from processed food.	500 g
TN1014	<b>ChroMedium MLGA (Membrane Lactose Glucuronide Agar)</b>	for 5.7 l culture medium (88.0 g for 1 l culture medium) Differentiation and enumeration of <i>E. coli</i> and other coliforms by a single membrane filtration technique.	500 g
TN1115	<b>CLED Agar (Brolacin Agar)</b>	for 14.5 l culture medium (34.3 g for 1 l culture medium) Determination of the colony count, isolation and initial differentiation of urinary tract pathogens.	500 g
TN1116	<b>Clostridial Differential Broth (DRCM)</b>	for 16.6 l culture medium: single strength (30.1 g for 1 l medium) for 8.3 l culture medium: double strength (60.2 g for 1 l medium) Determination of the colony count of <i>Clostridia</i> by means of the MPN method in foodstuffs and other materials.	500 g
TN1118	<b>Columbia Agar (Base) (Columbia Blood Agar Base)</b>	for 12.5 l culture medium (40.0 g for 1 l culture medium) Cultivation of fastidious microorganisms and to detect haemolysis. Addition of blood.	500 g
TN1191	<b>Columbia Agar acc. to harm. EP/USP/JIP</b>	for 12.2 l culture medium (41.0 g for 1 l culture medium) Test for <i>Clostridia</i> . Culture medium for testing non-sterile products. This culture medium is based on the recommendations of the harmonised method acc. to EP/USP/JIP (2006). Addition of blood.	500 g
TN1197	<b>Count Agar, sugar-free acc. to FIL-IDF</b>	for 14.3 l culture medium (35.0 g for 1 l culture medium) Determination of the germ count in milk and dairy products.	500 g

## D

Art. No.	Product	Description	Packing
TN1054	<b>Dermatophytes Agar, modified</b>	for 10.8 l culture medium (46.5 g for 1 l culture medium) Culture medium with Chloramphenicol and Cycloheximide included. Selective cultivation of dermatophytes and <i>Candida albicans</i> . No supplements necessary.	500 g
TN1121	<b>Desoxycholate Citrate Agar, modified (Leifson Agar, DCLS Agar, modified)</b>	for 10.1 l culture medium (49.5 g for 1 l culture medium) Detection and isolation of salmonellae and shigellae.	500 g
TN1124	<b>DEV Gelatin Agar</b>	for 11.1 l culture medium (45.0 g for 1 l culture medium) Used in water and waste water testing for the determination of the complete germ count and to detect gelatinase secreting cells.	500 g
TN1127 TN1127-01	<b>DEV Lactose Peptone Broth</b>	for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) Enrichment and determination of the germ count of <i>E. coli</i> and coliform bacteria in water.	500 g 5 kg
TN1128 TN1128-01	<b>DEV Nutrient Agar</b>	for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) Determination of complete germ counts in water acc. to DIN 38411 and § 64 LFGB German Food and Feed Code.	500 g 5 kg
TN1013	<b>DIASSALM (Diagnostic Semi-Solid Salmonella Agar)</b>	for 8.7 l culture medium (57.2 g for 1 l culture medium) Semi-solid culture medium for the detection of motile salmonellae from food samples and other test material.	500 g
TN1243 TN1243-01	<b>Double Buffered Peptone Water (Peptone Water Double Buffered)</b>	for 20.0 l culture medium (25.0 g for 1 l culture medium) for 200 l culture medium (25.0 g for 1 l culture medium) Pre-enrichment of pathogenic <i>Enterobacteriaceae</i> from foodstuffs and other test material, in particular with highly sour or acid-producing products.	500 g 5 kg

## E

Art. No.	Product	Description	Packing
TN1130	<b>Endo Agar (Lactose Fuchsin Sulfite Agar)</b>	for 14.5 l culture medium (34.6 g for 1 l culture medium) Detection and isolation of <i>Enterobacteriaceae</i> . No supplements necessary.	500 g
TN1236	<b>Enterobacteriaceae Enrichment Broth-Mossel acc. to harm. EP/USP/JP</b>	for 11.5 l culture medium (43.4 g for 1 l culture medium) Test for bile-tolerant gram-negative bacteria. Culture medium for testing non-sterile products. This culture medium is based on the recommendations of the harmonised method acc. to EP/USP/JP (2006).	500 g
TN1132	<b>Enterococci Selective Agar acc. to Slanetz-Bartley (Enterococci Agar)</b>	for 12.0 l culture medium (41.5 g for 1 l culture medium) Isolation and identification of enterococci especially from water (acc. to DIN EN ISO 7899-2) and from foodstuffs (§ 64 German Food and Feed Code).	500 g

# Dehydrated Culture Media

Art. No.	Product	Description	Packing
TN1200	<b>Eosin Methylene Blue Lactose Saccharose Agar (EMB Agar, Levine Agar)</b>	for 15.4 l culture medium (32.5 g for 1 l culture medium) Isolation of pathogenic <i>Enterobacteriaceae</i> , in particular salmonellae and shigellae, and for the cultivation and pre-differentiation of <i>Candida albicans</i> .	500 g

## F

Art. No.	Product	Description	Packing
TN1257	<b>Fraser Broth (Base)</b>	for 8.7 l culture medium (57.4 g for 1 l culture medium) Selective enrichment and the detection of <i>Listeria</i> spp. from foodstuffs (acc. to § 64 German Food and Feed Code) and from environmental materials. Addition of Fraser Selective Supplement TN 1318 or ½ Fraser Selective Supplement TN 1319.	500 g
TN1035	<b>Fraser Broth Complete</b>	for 8.6 l culture medium (57.9 g for 1 l culture medium) Selective enrichment of <i>Listeria</i> spp. from foodstuffs (acc. to § 64 German Food and Feed Code) and from environmental material. No supplements necessary.	500 g
TN1034 TN1034-01	<b>½ Fraser Broth Complete</b>	for 8.6 l culture medium (57.9 g for 1 l culture medium) for 86.3 l culture medium (57.9 g for 1 l culture medium) Selective enrichment of <i>Listeria</i> spp. from foodstuffs (acc. to § 64 German Food and Feed Code) and from environmental material. No supplements necessary.	500 g 5 kg

## G

Art. No.	Product	Description	Packing
TN1079	<b>Gelatin Lactose Medium</b>	for 3.2 l culture medium (155.0 g for 1 l culture medium) Confirmation of <i>Clostridium perfringens</i> (acc. to § 64 German Food and Feed Code).	500 g
TN1086	<b>Glucose Agar</b>	for 13.0 l culture medium (38.5 g for 1 l culture medium) Detection and colony count of <i>Enterobacteriaceae</i> in food.	500 g
TN1140	<b>Glucose Nutrient Broth</b>	for 17.5 l culture medium (28.5 g for 1 l culture medium) Used for the detection of aerobic bacteria, especially when testing for sterility.	500 g
TN1139	<b>Glucose Yeast Extract Cysteine Agar (Base) acc. to Beerens</b>	for 13.8 l culture medium (36.3 g for 1 l culture medium) Cultivation of anaerobes. Blood addition necessary.	500 g
TN1271	<b>GSP Agar (Base) (Glutamate Starch Phenol Red Agar, Pseudomonas Aeromonas acc. to Kielwein)</b>	for 11.1 l culture medium (44.9 g for 1 l culture medium) Detection of <i>Pseudomonas</i> sp. and <i>Aeromonas</i> sp. in all kinds of foodstuffs.	500 g



## H

Art. No.	Product	Description	Packing
TN1145	<b>HIB Medium (Base) (Urea Indole Motility Medium)</b>	for 5.4 l culture medium (28.0 g for 1 l culture medium) Differentiation of <i>Enterobacteriaceae</i> by means of detection of urea splitting, indole formation and motility. Addition of urea TN 1308.	150 g

## K

Art. No.	Product	Description	Packing
TN1217	<b>Kanamycin Esculin Azide Agar (KAA Agar)</b>	for 11.7 l culture medium (42.7 g for 1 l culture medium) Isolation, differentiation and determination of the colony count of enterococci from foodstuffs, water and other test material. No supplements necessary.	500 g
TN1218	<b>Kanamycin Esculin Azide Broth (KAA Broth)</b>	for 15.3 l culture medium (32.7 g for 1 l culture medium) Selective Broth for the cultivation of enterococci. No supplements necessary.	500 g
TN1287	<b>King's Agar (Base)</b>	for 2.7 l culture medium (36.9 g for 1 l culture medium) Detection and determination of the germ count of fluorescing bacteria in water, in particular of <i>Pseudomonas aeruginosa</i> (German Food and Feed Code). Addition of glycerin TN 1424.	100 g
TN1146	<b>Kligler Iron Agar (KIA, Double Sugar Iron Agar acc. to Kligler)</b>	for 10.1 l culture medium (49.6 g for 1 l culture medium) Differentiation of gram-negative bacteria, in particular <i>Enterobacteriaceae</i> . Urea TN 1308 may be added.	500 g

## L

Art. No.	Product	Description	Packing
TN1151	<b>Lactose Broth (Lactose Monohydrate Broth)</b>	for 38.5 l culture medium (13.0 g for 1 l culture medium) Cultivation and identification of coliform bacteria, in particular <i>E. coli</i> in water, food, pharmaceutical products and raw materials.	500 g
TN1278	<b>Lactose TTC Agar with Tergitol-7 (Tergitol-7 TTC Agar)</b>	for 9.4 l culture medium (53.2 g for 1 l culture medium) Isolation and differentiation of <i>E. coli</i> and coliform bacteria from water by means of the membrane filtration method. No supplements necessary.	500 g
TN1152	<b>Lauryl Sulfate Broth (Lauryl Sulfate Tryptose Lactose Broth)</b>	for 14.0 l culture medium (35.6 g for 1 l culture medium) Detection and determination of the germ count of coliform bacteria from water, waste water, foodstuffs and dairy products.	500 g
TN1153	<b>Lauryl Sulfate Tryptose Broth with Tryptophan and MUG (LST/MUG Medium)</b>	for 13.6 l culture medium: single strength (36.7 g for 1 l medium) for 6.8 l culture medium: double strength (73.4 g for 1 l medium) Detection and determination of the germ count of coliform bacteria from water, waste water, foodstuffs and dairy products.	500 g
TN1048	<b>LB Agar (Luria-Bertani Agar acc. to Miller)</b>	for 13.5 l culture medium (37.0 g for 1 l culture medium) Cultivation of <i>E. coli</i> in fermentation and molecular biology.	500 g

# Dehydrated Culture Media

Art. No.	Product	Description	Packing
TN1049	<b>LB Broth (Luria-Bertani Medium acc. to Miller)</b>	for 20.0 l culture medium (25.0 g for 1 l culture medium) Cultivation of <i>E. coli</i> in fermentation and molecular biology.	500 g
TN1066	<b>Linden Grain Medium</b>	for 847.5 l culture medium (29.5 g for 1 l culture medium) Cultivation of environmental germs, e.g. when testing beverage bottling installations.	25 kg
TN1154	<b>Lysine Decarboxylase Broth (LDC Broth)</b>	for 5.5 l culture medium (9.1 g for 1 l culture medium) Detection of lysine decarboxylase production by salmonellae and certain other <i>Enterobacteriaceae</i> .	50 g

## M

Art. No.	Product	Description	Packing
TN1156	<b>MacConkey Agar</b>	for 10.3 l culture medium (48.5 g for 1 l culture medium) Isolation of <i>Enterobacteriaceae</i> from stool, urine, food, and waste water samples and other materials, also in cases involving a high incidence of salmonellae and shigellae.	500 g
TN1075 TN1075-01	<b>MacConkey Agar acc. to harm. EP/USP/JPP</b>	for 10.0 l culture medium (50.0 g for 1 l culture medium) for 100.0 l culture medium (50.0 g for 1 l culture medium) Test for <i>Escherichia coli</i> . Culture medium for testing nonsterile products. This culture medium is based on the recommendations of the harmonised method acc. to EP/USP/JPP (2006).	500 g 5 kg
TN1027	<b>MacConkey Broth acc. to harm. EP/USP/JPP</b>	for 14.3 l culture medium (35.0 g for 1 l culture medium) Test for <i>Escherichia coli</i> . Culture medium for testing nonsterile products. This culture medium is based on the recommendations of the harmonised method acc. to EP/USP/JPP (2006).	500 g
TN1157	<b>Magnesium Chloride Malachite Green Broth acc. to Rappaport-Vassiliadis (RV Medium)</b>	for 15.0 l culture medium (33.4 g for 1 l culture medium) Used for selective enrichment during isolation of <i>salmonellae</i> from foodstuffs and environmental materials.	500 g
TN1158	<b>Malachite-green Broth</b>	for 56.8 l culture medium (8.8 g for 1 l culture medium) Detection of <i>Pseudomonas aeruginosa</i> from water (acc. to § 64 German Food and Feed Code). No supplements necessary.	500 g
TN1267	<b>Malt Extract Agar (Wort Peptone Agar)</b>	for 10.0 l culture medium (50.0 g for 1 l culture medium) Detection, isolation and determination of the germ count of yeasts and fungi in foodstuffs and other test materials.	500 g

Art. No.	Product	Description	Packing
TN1166	<b>Mannitol Salt Agar acc. to harm. EP/USP/JP</b>	for 4.5 l culture medium (111.0 g for 1 l culture medium) Test for <i>Staphylococcus aureus</i> . Culture medium for testing non-sterile products. This culture medium is based on the recommendations of the harmonised method acc. to EP/USP/JP (2006).	500 g
TN1288	<b>m-CP Agar (Base) (Chromogenic Membrane Filtration Clostridium perfringens Agar)</b>	for 7.3 l culture medium (68.1 g for 1 l culture medium) Used to detect <i>Clostridium perfringens</i> (including spores) in water. Addition of m-CP-Selective Supplement TN 1330.	500 g
TN1161	<b>Methyl-red Voges Proskauer Broth (MRVP)</b>	for 8.8 l culture medium (17.0 g for 1 l culture medium) Used to differentiate the <i>coli</i> -aerogenes group by means of the methyl red test and the Voges-Proskauer reaction.	150 g
TN1040	<b>Minca Agar, modified</b>	for 18.9 l culture medium (26.4 g for 1 l culture medium) Cultivation of <i>E. coli</i> relevant to veterinary medicine (young cattles). Addition of Minca Supplement TN 1334.	500 g
TN1071	<b>Modified Scholtens' Broth (MSB)</b>	for 17.0 l culture medium (29.4 g for 1 l culture medium) Used to detect and enumerate somatic coliphages in all kinds of water, sediments and sludges (DIN EN ISO 10705-2).	500 g
TN1041	<b>Modified Soybean Casein Digest Broth (Mod. Tryptic Soy Broth, m-TSB)</b>	for 15.2 l culture medium (33.0 g for 1 l culture medium) Selective enrichment of <i>E. coli</i> O 157 from food samples (§ 64 German Food and Feed Code) and from faeces acc. to DIN EN ISO 16654. Novobiocin Selective Supplement may be added.	500 g
TN1201	<b>MRS Agar (Base) (Lactobacillus Agar acc. to de Man, Rogosa and Sharpe)</b>	for 8.0 l culture medium (62.3 g for 1 l culture medium) Enrichment, cultivation and determination of the germ count of all <i>Lactobacillus</i> species from meat, milk and dairy products (acc. to § 64 German Food and Feed Code) and from other test materials. Addition of Tween® 80 TN 1422.	500 g
TN1068	<b>MRS Agar (Base) with pH 5.7</b>	for 7.8 l culture medium (64.3 g for 1 l culture medium) Enrichment, cultivation and determination of the germ count of all <i>Lactobacillus</i> species from meat, milk and dairy products (acc. to § 64 German Food and Feed Code) and from other test materials. Addition of Tween 80® TN 1422.	500 g
TN1205	<b>MRS Broth (Base)</b>	for 9.7 l culture medium (51.3 g for 1 l culture medium) Enrichment, cultivation and determination of the germ count of all <i>Lactobacillus</i> species from meat, milk and dairy products (acc. to § 64 German Food and Feed Code) and from other test materials. Addition of Tween® 80 TN 1422.	500 g
TN1272	<b>MSRV Medium Base (Modified Semi-Solid Rappaport Vassiliadis Medium Base)</b>	for 15.8 l culture medium (31.7 g for 1 l culture medium) Semi-solid culture medium for the detection of motile salmonellae from food samples and other test material. Addition of Novobiocin Selective Supplement TN 1324.	500 g
TN1162	<b>Mueller Hinton Agar (MHA)</b>	for 13.2 l culture medium (38.0 g for 1 l culture medium)	500 g
TN1250		for 131.6 l culture medium (38.0 g for 1 l culture medium) For use in the antimicrobial sensitivity test using the agar diffusion method.	5 kg

# Dehydrated Culture Media

Art. No.	Product	Description	Packing
TN1163	<b>Mueller Hinton Broth (MHB)</b>	for 23.8 l culture medium (21.0 g for 1 l culture medium) Used to perform the sensitivity test according to the macrodilution and microdilution methods (MIC test).	500 g
TN1291	<b>MYP Agar (Base) (Cereus Selective Agar Base acc. to Mossel)</b>	for 11.6 l culture medium (43.0 g for 1 l culture medium) Determination of the colony count, detection and isolation of <i>Bacillus cereus</i> from foodstuffs (acc. to § 64 German Food and Feed Code). Addition of Egg Yolk Emulsion TN 1316 and Bacillus Cereus Selective Supplement TN 1315.	500 g

## N

Art. No.	Product	Description	Packing
TN1164 TN1164-01	<b>Nutrient Agar I</b>	for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) Universal culture medium for the cultivation of fastidious microorganisms. Blood may be added.	500 g 5 kg
TN1168	<b>Nutrient Agar III</b>	for 26.3 l culture medium (19.0 g for 1 l culture medium) Cultivation of suspicious bacteria from test materials for subsequent use in biochemical or serological identification.	500 g
TN1176	<b>Nutrient Agar acc. to the German drinking water regulation and Food and Feed Code</b>	for 20 l culture medium (25.0 g for 1 l culture medium) Universal culture medium for cultivation of fastidious microorganisms (acc. to the German drinking water regulation (TrinkwV) and German Food and Feed Code (§ 64 LFGB)). Culture medium is composed in accordance with DIN EN 16266. Blood may be added.	500 g
TN1171 TN1171-01	<b>Nutrient Broth</b>	for 62.5 l culture medium (8.0 g for 1 l culture medium) for 625 l culture medium (8.0 g for 1 l culture medium) Cultivation of less fastidious microorganisms.	500 g 5 kg
TN1172	<b>Nutrient Broth I</b>	for 20 l culture medium (25.0 g for 1 l culture medium) Nutrient-rich culture medium for cultivation of microorganisms, including fastidious microorganisms.	500 g
TN1174	<b>Nutrient Broth II</b>	for 33.3 l culture medium (15.0 g for 1 l culture medium) Cultivation of microorganisms.	500 g

## O

Art. No.	Product	Description	Packing
TN1207	<b>Orange Serum Agar</b>	for 12.5 l culture medium (40.0 g for 1 l culture medium) Isolation, cultivation and determination of the germ count of acid-tolerant spoilage organisms in fruit juices and fruit extracts.	500 g

Art. No.	Product	Description	Packing
TN1269	<b>Orange Serum Broth</b>	for 20 l culture medium (25.0 g for 1 l culture medium) Isolation, cultivation and determination of the germ count of acid-tolerant spoilage organisms in fruit juices and fruit extracts.	500 g

## P

Art. No.	Product	Description	Packing
TN1209	<b>Palcam Agar (Base) (Palcam Agar acc. to van Netten)</b>	for 7.4 l culture medium (68.0 g for 1 l culture medium) Isolation and detection of <i>Listeria monocytogenes</i> from foodstuffs (acc. to § 64 German Food and Feed Code), biological samples and strongly contaminated samples from the environment. Addition of Palcam Selective Supplement TN 1312.	500 g
TN1147	<b>Peptone Saline Diluent (Maximum Recovery Diluent)</b>	for 52.6 l culture medium (9.5 g for 1 l culture medium) Culture medium for preparing samples acc. to DIN EN ISO 6887-1.	500 g
TN1189	<b>Plate Count Agar (TGE Agar, Tryptone Glucose Yeast Extract Agar)</b>	for 27.0 l culture medium (18.5 g for 1 l culture medium) Used to determine complete germ counts in milk, dairy products (acc. to § 64 German Food and Feed Code), water and other test materials.	500 g
TN1190	<b>Plate Count Agar with Skimmed Milk (Tryptone Glucose Yeast Extract Milk Agar)</b>	for 25.6 l culture medium (19.5 g for 1 l culture medium) Determination of germ counts in milk and dairy products (acc. to § 64 German Food and Feed Code).	500 g
TN1150	<b>Potato Dextrose Agar acc. to harm. EP/USP/JP</b>	for 12.8 l culture medium (39.0 g for 1 l culture medium) Preparation of <i>Aspergillus niger</i> test strain and for isolation. Used to cultivate and to maintain strains of yeasts and fungi. Medium for testing non-sterile products based on the recommendations of the harmonised method acc. to EP/USP/JP (2006).	500 g
TN1286	<b>Pseudomonas Agar (Base)</b>	for 10.3 l culture medium (48.4 g for 1 l culture medium) Used to detect and count <i>Pseudomonas aeruginosa</i> from water samples, foodstuffs and other material. Addition of Pseudomonas CN Selective Supplement TN 1323 and glycerin TN 1424.	500 g

## R

Art. No.	Product	Description	Packing
TN1297	<b>R2A Agar</b>	for 27.6 l culture medium (18.1 g for 1 l culture medium) Used in the testing of water for use in injections.	500 g
TN1262	<b>Rappaport Vassiliadis Salmonella Enrichment Broth acc. to harm. EP/USP/JP (RVS Broth)</b>	for 18.5 l culture medium (27.1 g for 1 l culture medium) Selective enrichment of salmonellae. This culture medium is based on the recommendations of the harmonised method acc. to EP/USP/JP (2006).	500 g

# Dehydrated Culture Media

Art. No.	Product	Description	Packing
TN1167	<b>Reinforced Medium for Clostridia acc. to harm. EP/USP/JP (RCM)</b>	for 13.2 l culture medium (38.0 g for 1 l culture medium) Test for <i>Clostridia</i> . Selective enrichment of salmonellae. This culture medium is based on the recommendations of the harmonised method acc. to EP/USP/JP (2006).	500 g
TN1039	<b>Rice Extract Agar</b>	for 29.4 l culture medium (17.0 g for 1 l culture medium) Differentiation of <i>Candida albicans</i> via the formation of chlamydo spores possible. Addition of Tween® 80 TN 1422.	500 g

## S

Art. No.	Product	Description	Packing
TN1264	<b>Sabouraud Dextrose Agar acc. to harm. EP/USP/JP</b>	for 7.7 l culture medium (65.0 g for 1 l culture medium) Test for <i>Candida albicans</i> . Culture medium for testing non-sterile products. Culture medium is based on the recommendations of the harmonised method acc. to EP/USP/JP (2006).	500 g
TN1047	<b>Sabouraud Dextrose Broth acc. to harm. EP/USP/JP</b>	for 16.7 l culture medium (30.0 g for 1 l culture medium) Test for <i>Candida albicans</i> . Culture medium for testing non-sterile products. Culture medium is based on the recommendations of the harmonised method acc. to EP/USP/JP (2006).	500 g
TN1178 TN1178-01	<b>Sabouraud 2%Glucose Agar (Sabouraud Dextrose Agar)</b>	for 11.9 l culture medium (42.0 g for 1 l culture medium) for 119.0 l culture medium (42.0 g for 1 l culture medium) Used to cultivate and maintain strains of dermatophytes, yeasts and fungi.	500 g 5 kg
TN1016	<b>Sabouraud 2%Glucose Chloramphenicol Agar</b>	for 11.9 l culture medium (42.1 g for 1 l culture medium) Used to isolate, cultivate and maintain strains of pathogenic and apathogenic dermatophytes, yeasts and moulds.	500 g
TN1015	<b>Sabouraud Glucose Medium with Chloramphenicol</b>	for 8.1 l culture medium (62.1 g for 1 l culture medium) Selective cultivation of fungi in sterility testing.	500 g
TN1181	<b>Schaedler Agar (Base)</b>	for 13.4 l culture medium (37.2 g for 1 l culture medium) Cultivation and application of the antimicrobial sensitivity test to anaerobes.	500 g
TN1203	<b>Schaedler Broth</b>	for 19.1 l culture medium (26.2 g for 1 l culture medium) Cultivation and application of the antimicrobial sensitivity test to anaerobes.	500 g
TN1225	<b>Selenite Broth</b>	for 21.7 l culture medium (23.0 g for 1 l culture medium) Selective enrichment of salmonellae from faeces, foodstuffs and other materials. No supplements necessary.	500 g
TN1183	<b>Selenite Cystine Broth</b>	for 21.7 l culture medium (23.0 g for 1 l culture medium) Selective enrichment of salmonellae from faeces, foodstuffs and other test materials. No supplements necessary.	500 g
TN1184	<b>Simmons Citrate Agar</b>	for 9.8 l culture medium (15.3 g for 1 l culture medium) Differentiation of <i>Enterobacteriaceae</i> on the basis of the utilisation of citrate as the sole source of carbon (acc. to DIN 38411, § 64 German Food and Feed Code).	150 g

Art. No.	Product	Description	Packing
TN1220	<b>Sorbitol MacConkey Agar (SMAC Agar)</b>	for 10.3 l culture medium (48.5 g for 1 l culture medium) Cultivation and pre-differentiation of <i>E. coli</i> serogroup O 157.	500 g
TN1221 TN1221-01	<b>SS Agar (Salmonella Shigella Agar Modified)</b>	for 8.8 l culture medium (56.5 g for 1 l culture medium) for 88.5 l culture medium (56.5 g for 1 l culture medium) Isolation of salmonellae and shigellae. Inhibits the growth of gram-positive microorganisms, suppresses swarming of <i>Proteus</i> sp.	500 g 5 kg
TN1017	<b>Synthetic Nutrient Deficient Agar (SNA)</b>	for 29.1 l culture medium (17.2 g for 1 l culture medium) Cultivation of yeasts and moulds in particular in plant culture.	500 g

## T

Art. No.	Product	Description	Packing
TN1259	<b>TBX Chromogen Agar (Tryptone Bile X Glucuronic Agar)</b>	for 15.8 l culture medium (31.6 g for 1 l culture medium) Detection and determination of the colony count of <i>E. coli</i> from food. No supplements necessary.	500 g
TN1187	<b>Tetrathionate Broth (Base) acc. to Muller-Kauffmann</b>	for 5.0 l culture medium (99.9 g for 1 l culture medium) Selective enrichment of salmonellae from meat, meat products and other foodstuffs. Addition of iodine-potassium iodide solution (5 g potassium iodide, 4 g iodine, 20 ml distilled water).	500 g
TN1188 TN1188-01	<b>Thioglycolate Medium acc. to EP/USP (Fluid Thioglycolate Medium)</b>	for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) Detection of aerobe and anaerobe microorganisms and when testing the sterility of products (EP and USP).	500 g 5 kg
TN1256	<b>Thioglycolate Medium Modified (high degree of transparency)</b>	for 16.7 l culture medium (30.0 g for 1 l culture medium) Detection of aerobe and anaerobe microorganisms and when testing the sterility of products.	500 g
TN1023	<b>Thioglycolate Medium without Indicator</b>	for 16.6 l culture medium (30.1 g for 1 l culture medium) Detection of a variety of microorganisms, in particular anaerobes, from clinical material.	500 g
TN1083 TN1083-01	<b>Tryptic Soy (CASO) Broth Irradiated (Soyabean Casein Digest Broth Irradiated)</b>	for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) Sterility control of filling equipment (media-fill).	500 g 5 kg
TN1260	<b>Tryptone Bile Agar (TBA)</b>	for 15.9 l culture medium (31.5 g for 1 l culture medium) Used to detect and count <i>E. coli</i> using the membrane filtration method in water and the membrane-agar method in food (acc. to § 64 German Food and Feed Code).	500 g

# Dehydrated Culture Media

Art. No.	Product	Description	Packing
TN1036	<b>Tryptone Soy Yeast Extract Agar (TSYEA)</b>	for 10.4 l culture medium (48.0 g for 1 l culture medium) Confirmation of <i>Listeria</i> spp. from milk and other dairy products and other foodstuffs.	500 g
TN1258	<b>Tryptophan Pepton Water (Tryptophan Broth)</b>	for 6.3 l culture medium (16.0 g for 1 l culture medium) Differentiation of <i>Enterobacteriaceae</i> by detection of indole formation (DIN EN 9308-1).	100 g
TN1241	<b>TSC Agar (Base) (Tryptose Sulfit Cycloserine Agar)</b>	for 12.8 l culture medium (39.0 g for 1 l culture medium) Detection of <i>Clostridium perfringens</i> and other sulfitereducing <i>Clostridia</i> taken from foodstuffs and water. Addition of TSC Selective Supplement TN 1333.	500 g

## U

Art. No.	Product	Description	Packing
TN1143	<b>Urea Agar (Base) acc. to Christensen</b>	for 14.7 l culture medium (17.0 g for 1 l culture medium) Used for the urease-test for the presence of urease-positive bacteria, e.g. <i>Proteus</i> sp. and <i>Klebsiella</i> sp. (acc. to § 64 German Food and Feed Code).	250 g

## V

Art. No.	Product	Description	Packing
TN1148	<b>Violet Red Bile Glucose Agar (VRBD Agar)</b>	for 13.3 l culture medium (37.5 g for 1 l culture medium) Isolation and determination of the germ count of <i>Enterobacteriaceae</i> in foodstuffs.	500 g
TN1042	<b>Violet Red Bile Glucose Agar acc. to harm. EP/USP/JP (VRBD Agar)</b>	for 13.7 l culture medium (36.5 g for 1 l culture medium) Test for bile-tolerant gram-negative bacteria. Culture medium for testing non-sterile products. This culture medium is based on the recommendations of the harmonised method acc. to EP/USP/JP (2006).	500 g
TN1149	<b>Violet Red Bile Lactose Agar (VRBL Agar, VRB Agar)</b>	for 13.3 l culture medium (37.5 g for 1 l culture medium) Detection and determination of the colony count of coliform bacteria from water, foodstuffs and dairy products.	500 g



## W

Art. No.	Product	Description	Packing
TN1194 TN1194-01	<b>Water-blue Metachrome-yellow Lactose Agar acc. to Gassner, Modified (Gassner Agar, mod.)</b>	for 9.9 l culture medium (50.3 g for 1 l culture medium) for 99.4 l culture medium (50.3 g for 1 l culture medium) Detection and isolation of <i>Enterobacteriaceae</i> from foodstuffs (especially meat) and other test materials.	500 g 5 kg
TN1210	<b>Wort Agar (Base)</b>	for 10.4 l culture medium (48.0 g for 1 l culture medium) Detection, isolation and determination of the colony count of yeasts and fungi (especially in controls carried out during the production of beverages). Addition of glycerin TN 1424.	500 g
TN1211	<b>Wort Broth (Base)</b>	for 15.2 l culture medium (33.0 g for 1 l culture medium) Detection, isolation and determination of the colony count of yeasts and fungi. Addition of glycerin TN 1424.	500 g

## X

Art. No.	Product	Description	Packing
TN1170	<b>XLD Agar acc. to harm. EP/USP/JP (Xylose Lysine Deoxycholate Agar)</b>	for 9.1 l culture medium (55.2 g for 1 l culture medium) Test for <i>Salmonella</i> . Medium for testing non-sterile products. Medium is based on the recommendation of the harmonised method acc. to EP/USP/JP (2006).	500 g
TN1196	<b>XLD Agar acc. to ISO 6579 (Xylose Lysine Desoxycholate Agar)</b>	for 8.8 l culture medium (57.0 g for 1 l culture medium) Isolation of salmonellae and shigellae from foodstuffs, clinical materials and pharmaceuticals.	500 g

## Y

Art. No.	Product	Description	Packing
TN1290	<b>Yeast Extract Agar</b>	for 26.3 l culture medium (19.0 g for 1 l culture medium) Used to determine the colony count of microorganisms cultivable at 22 °C and 35 °C from water and waste water (DIN EN ISO 6222).	500 g
TN1144	<b>Yeast Extract Glucose Chloramphenicol Agar (YGC Agar)</b>	for 14.2 l culture medium (35.1 g for 1 l culture medium) Detection and determination of the colony count of yeasts and moulds from foodstuffs. No supplements necessary.	500 g
TN1229	<b>Yersinia Agar (Base) acc. to Schiemann (CIN)</b>	for 8.9 l culture medium (56.0 g for 1 l culture medium) Isolation of <i>Yersinia enterocolitica</i> from foodstuffs and clinical test materials. Addition of Yersinia Selective Supplement TN 1301.	500 g

# Supplements

## Supplements

Art. No.	Product	Description	Packing
TN1315	<b>Bacillus Cereus Selective Supplement (Polymyxin B Selective Supplement)</b>	for 6 l culture medium (2 vials for 1 culture medium) Inhibition of accompanying flora during the selection of <i>Bacillus cereus</i> . In conjunction with Bacillus Cereus Agar (Base) TN 1245 and Egg Yolk Emulsion TN 1316.	12 x 1 vial
TN1304	<b>Campylobacter Growth Supplement</b>	for 22.2 l culture medium (4.5 g for 1 l culture medium) Used to promote growth during the cultivation of <i>Campylobacter</i> spp. In conjunction with Nutrient Agar I TN 1164, TN 1165 and potato starch.	100 g
TN1302	<b>Campylobacter Selective Supplement</b>	for 6 l culture medium (2 vials for 1 l culture medium) Suppression of accompanying faecal flora during the cultivation of <i>Campylobacter</i> spp., in particular <i>Campylobacter fetus</i> . In conjunction with Campylobacter Growth Supplement TN 1304, Nutrient Agar I TN 1164, TN 1165 and potato starch.	12 x 1 vial
TN1316	<b>Egg Yolk Emulsion</b>	for approx. 2 l culture medium (2 vials for 1 l culture medium) Isolation and differentiation of <i>Bacillus</i> spp., in particular of <i>Bacillus cereus</i> from foodstuffs (acc. to § 64 German Food and Feed Code). Supplement for culture media: 1 vial contains 25 ml 2 vials for 950 ml Bacillus C. Agar (Base) TN 1245. 4 vials for 900 ml MYP Agar (Base) TN 1291.	4 x 25 ml
TN1310	<b>Egg Yolk Potassium Tellurite Emulsion</b>	for approx. 2 l culture medium (2 vials for 1 l culture medium) Isolation and differentiation of <i>Staphylococcus aureus</i> from foodstuffs and pharmaceutical products. Supplement for culture media: 1 vial contains 25 ml 2 vials for 950 ml Baird Parker Agar (Base) TN 1104.	4 x 25 ml
TN1318	<b>Fraser Selective Supplement</b>	for 3 l culture medium (1 vial per component for 0.5 l medium) Inhibition of accompanying flora during the selective enrichment of <i>Listeria monocytogenes</i> . Supplement for culture media consists of two components. 1 vial per component for 0.5 l Fraser Broth (Base) TN 1257.	6 x 2 vials
TN1319	<b>½ Fraser Selective Supplement</b>	for 13.5 l culture medium (1 vial per component for 2.25 l medium) Inhibition of accompanying flora during the selective enrichment of <i>Listeria monocytogenes</i> . Supplement for culture media consists of two components. 1 vial per component for 2.25 l Fraser Broth (Base) TN 1257.	6 x 2 vials
TN1330	<b>m-CP Selective Supplement</b>	for 2 l culture medium (1 vial per component for 0.5 l culture medium) Used to detect <i>Clostridium perfringens</i> (incl. spores) in water. Supplement for culture media consists of three components. 1 vial per component for 0.5 l m-CP Agar (Base) TN 1288.	4 x 3 vials

Art. No.	Product	Description	Packing
TN1334	<b>Minca Supplement</b>	for 1.2 l culture medium (1 vial for 0.1 l culture medium) Encourage development of fimbrial antigen F 5 (K 99) in enteropathogenic <i>E. coli</i> . Relevant to veterinary medicine (young cattles). 1 vial for 100 ml Minca Agar, modified TN 1040.	12 x 1 vials
TN1324	<b>Novobiocin Selective Supplement</b>	for 6 l culture medium (2 vials for 1 l culture medium) Detection of motile salmonellae from foodstuffs and other test materials. 1 vial for 0.5 l MSRV Medium (Base) TN 1272.	12 x 1 vial
TN1312	<b>Palcam Selective Supplement</b>	for 6 l culture medium (2 vials for 1 l culture medium) Inhibition of accompanying flora during the selection of <i>Listeria monocytogenes</i> . 1 vial for 0.5 l Palcam Agar (Base) TN 1209.	12 x 1 vial
TN1323	<b>Pseudomonas CN Selective Supplement</b>	for 6 l culture medium (2 vials for 1 l culture medium) Inhibition of accompanying flora during the isolation of <i>Pseudomonas aeruginosa</i> . 1 vial for 0.5 l Pseudomonas Agar (Base) TN 1286.	12 x 1 vial
TN1333	<b>TSC Selective Supplement</b>	for 6 l culture medium (2 vials for 1 l culture medium) Inhibition of accompanying flora during the isolation of <i>Clostridium perfringens</i> . 1 vial for 0.5 l TSC Agar (Base) TN 1241.	12 x 1 vial
TN1301	<b>Yersinia Selective Supplement</b>	for 6 l culture medium (2 vials for 1 l culture medium) Inhibition of accompanying flora during the isolation of <i>Yersinia enterocolitica</i> . 1 vial for 0.5 l Yersinia Agar (Base) acc. to Schiemann (CIN) TN 1229.	12 x 1 vial

# Additives Detection Reagents

## Additives

Art. No.	Product	Description	Packing
TN1424	<b>Glycerin, water-free</b>	Carbon source and energy source, inhibits the growth of bacterial accompanying flora. Supplement for culture media: Cetrimide Agar acc. to harm. EP/USP/JP TN 1024 Bile Chrysoidine Glycerol Medium (Base) with MUG TN 1237 Bile Chrysoidine Glycerol Agar (Base) TN 1133, TN 1249 Buffered Nitrate Motility Medium TN 1078 King's Agar (Base) acc. to EN 12780 TN 1287 Pseudomonas Agar (Base) TN 1286 Wort Agar (Base) TN 1210 Wort Broth (Base) TN 1211	100 ml
TN1422	<b>Tween® 80</b>	Emulsifier for lipids and oils. Solubilizers for substances that are difficult to wet. Growth factor. Supplement for culture media: 1 ml for 1 l CATC Agar (Base) TN 1273. 1 ml for 1 l MRS Agar (Base) TN 1201. 1 ml for 1 l MRS Agar (Base) with pH 5.7 TN 1068. 1 ml for 1 l MRS Broth (Base) TN 1205. 10 ml for 1 l Rice Extract Agar TN 1039.	100 ml
TN1308	<b>Urea Solution, 40%</b>	Urea test for sterile culture media. Supplement for culture medium: 1 vial for 100 ml Urea Agar (Base) acc. to Christensen TN 1143. 1 vial for 100 ml HIB Medium (Base) TN 1145. 1 vial for 1 l Kligler Iron Agar TN 1146.	12 x 1 vial

## Detection reagents

Art. No.	Product	Description	Packing
TN1512	<b>Acetamide Solution</b>	for 20 tests	20 x 1 btl
TN1512-01		for 100 tests Used to detect ammonia in the identification of <i>Pseudomonas aeruginosa</i> . Ready-to-use Acetamide Solution acc. to DIN EN ISO 16266. 1 bottle contains 5 ml Acetamide Solution. Addition of Nessler's Reagent TN 1515, TN 1516.	100 x 1 btl
TN1519	<b>Acid Phosphatase Reagent</b>	Reagent is used in accordance with ISO 14189 to detect acid phosphatase from <i>Clostridium perfringens</i> . 6 vials dye reagent, 1 bottle buffer	6 vials + 1 btl
TN1515	<b>Nessler's Reagent</b>	for 60 vials Acetamide Solution	1 x 5 ml
TN1515-01		for 480 vials Acetamide Solution Used to detect ammonia in the identification of <i>Pseudomonas aeruginosa</i> . In conjunction with Acetamide Solution TN 1512, TN 1513.	4 x 10 ml
TN1517	<b>Oxidase Reagent</b>	for 12 tests Used for the oxidase reaction. 1 vial in 2 ml distilled water.	12 x 1 vial

# Ready-to-use Culture Media

## Ready-to-use culture media

Art. No.	Product	Description	Packing
TN1712	<b>Sabouraud Glucose Slant Agar</b>	Used to isolate, cultivate and maintain strains of pathogenic and apathogenic dermatophytes, yeasts and moulds. 1 tube contains 5 ml culture medium.	20 tubes
TN1714	<b>Sabouraud Glucose Slant Agar with Chloramphenicol and Cycloheximide</b>	Isolation of dermatophytes. 1 tube contains 5 ml culture medium.	20 tubes
TN1702	<b>Swarm Agar</b>	Enables salmonellae to swarm well, which promotes improved formation of the H-antigens. 1 tube contains 10 ml culture medium.	20 tubes

## Ready-to-use culture media for breweries

Art. No.	Product	Description	Packing
TN1724	<b>DEV Nutrient Agar</b>	Determination of complete germ counts in water acc. to DIN 38411 and § 64 LFGB (German Food and Feed Code).	9 x 250 ml
TN1743	<b>Lysine Agar</b>	Isolation and determination of the germ count of non-Saccharomyces (wild) yeasts in breweries.	9 x 250 ml
TN1707 TN1707-01	<b>MegaPec Broth</b>	Selective enrichment of gram-negative beer spoiling bacteria belonging to the genera <i>Pectinatus</i> and <i>Megasphaera</i> . 1 tube contains 12 ml broth.	20 tubes 100 tubes
TN1502	<b>VLB-S7-S Agar</b>	Detection of gram-positive, beer spoiling microorganisms (lactobacilli and pediococci in beer).	250 ml
TN1503	<b>VLB-S7-S Broth</b>	Detection of gram-positive, beer spoiling microorganisms (lactobacilli and pediococci in beer).	250 ml
TN1744	<b>WL Nutrient Agar (WLN)</b>	Used for the non-selective determination of microbial flora in breweries.	9 x 250 ml
TN1745	<b>WL Nutrient Agar with Cycloheximide (WLD)</b>	Detection and determination of the colony count of bacteria from test material that contain large numbers of yeasts and fungi.	9 x 250 ml
TN1746	<b>Wort Agar</b>	Cultivation of yeasts and moulds.	9 x 250 ml

# Base Materials

## Base materials

Art. No.	Product	Description	Packing
TN1400	<b>Agar for microbiological purpose</b>	Basal medium for the preparation of culture media for microbiological use.	500 g
TN1403 TN1403-01	<b>Casein peptone, pancreatic digested (Tryptone)</b>	Growth of a wide variety of organisms. Nutritious peptone and amino-acids. Component of many culture media.	500 g 5 kg
TN1423	<b>Glucose for microbiological use</b>	Dextrose for microbiological use.	500 g
TN1418	<b>Lactose for microbiology</b>	Milk sugar for microbiological use.	500 g
TN1410	<b>Meat extract</b>	Culture medium additive.	500 g
TN1405	<b>Meat peptone, pancreatic digested</b>	A highly nutritious peptone.	500 g
TN1417	<b>Sodium chloride for microbiological purpose</b>	Table salt for microbiological use.	1 kg
TN1419	<b>Soy bean Peptone, papainic digested</b>	A broad nutrient spectrum, a high level of carbohydrates and vitamins.	500 g
TN1404	<b>Yeast extract for microbiological use</b>	A source of carbon and amino acid and a rich source of B vitamins and growth factors.	500 g

Do you have a specific request?

We develop and produce according to your demand.

# Immunoassays

DIAGNOSTICS WITH PASSION



## Detection of rabies virus

Our Monoclonal Anti-Rabies, FITC, is used for detection of rabies virus in impression smears from different sources. The direct immunofluorescence test can be used for the detection of rabies

virus in brain tissues of animals and by virus isolation in cell cultures.

The product contains fluorescein isothiocyanate (FITC)-labelled monoclonal antibodies.

## Our portfolio comprises:

- Monoclonal Anti-Rabies, FITC



# Immunoassays

## Detection of rabies virus

Monoclonal Anti-Rabies, FITC is used for the detection of rabies virus in impression smears from brain of animals and by virus isolation in cell cultures. The test is based on the principle of direct immunofluorescence. The test samples are coated with Monoclonal Anti-Rabies, FITC in the recommended working dilution, washed and placed in a fluorescence microscope to test for fluorescence. This fluorescent antibody test (FAT) is the most widely used method for diagnosing rabies infection and recommended by both WHO and OIE.

Monoclonal Anti-Rabies, FITC contains fluorescein isothiocyanate (FITC)-conjugated monoclonal antibodies, which are isolated from cell culture supernatants and conjugated with FITC.

The product is approved by the Licensing Authority of FLI, Federal Research Institute for Animal Health.

License No.: FLI B-555

Every manufactured product batch is released by the OIE and National Reference Laboratory of FLI.

Art. No.	Product	Description	Packing
PA1202	<b>Monoclonal Anti-Rabies, FITC</b>	Sensitive detection of the classical rabies virus or bat associated lyssaviruses, especially EBLV-1 and EBLV-2 using the fluorescent antibody test (FAT).	1 ml



# Monoclonal Antibodies

DIAGNOSTICS WITH PASSION



## The perfect combination of technology and antibody

Use your own technology combined with our antibodies. We offer monoclonal antibodies for research or further use in your own technique.

Depending on your demand, we supply supernatant or purified antibodies. In case you need a special labeling for your application, just contact us.

## Our portfolio comprises:

- STEC-(VTEC) diagnostics
- Capture antibody for human IgM EIA
- Specific antibodies against *Salmonella*
- Anti-Mistletoe lectins for determining mistletoe lectin



Optimised management of your cell line and reliable production of antibodies in high concentrations

# Monoclonal Antibodies

## Anti-Human IgM

Art. No.	Product	Description	Packing
PA1312	<b>Anti-Human IgM</b>	capture antibody Sufficient for coating 10 micro-titre plates Coating antibody used for human IgM antibody detection by capture enzyme immunoassays.	1 mg

## Anti-Mistletoe lectins (A chain)

Art. No.	Product	Description	Packing
PL110008	<b>Anti-Mistletoe lectin (A chain) 5F5</b>	5F5 binds preferentially to native epitopes of mistletoe lectins I-III. Serves as capture antibody in ELISA in combination with detection antibody „anti-mistletoe lectin (A chain) 5H8, POD labelled“ for determining total lectin content in mistletoe extracts.	0.1 mg
PL110005	<b>Anti-Mistletoe lectin (A chain) 5H8, POD labelled</b>	5H8 binds preferentially to native epitopes of mistletoe lectins I-III. Serves as detection antibody in ELISA in combination with capture antibody „anti-mistletoe lectin (A chain) 5F5“ for determining total lectin content in mistletoe extracts.	0.1 ml

## Anti-Shigatoxin (Verotoxin)

Art. No.	Product	Description	Packing
TS2106	<b>Anti-Shigatoxin 1</b>	clone: VT 109/4-E9b Reacts with subunit B of Shigatoxin 1	1 mg
TS2103	<b>Anti-Shigatoxin 2</b>	clone: VT 135/6-B9 Reacts with subunit A of Shigatoxin 2	1 mg
TS2107	<b>Anti-Shigatoxin 2</b>	clone: VT 136/8-H4 Reacts with subunit B of Shigatoxin 2	1 mg

## Anti-Salmonella

Product	Packing
<b>Salmonella or Shigella specific antibodies</b>	on request

**TAILORED  
PRODUCTS**

„sifin diagnostics gmbh addresses all IVD-regulated markets in contract manufacturing and is among the few FDA-audited contract manufacturing companies producing monoclonal antibodies for blood grouping diagnostics.“

# Contract Manufacturing

DIAGNOSTICS WITH PASSION



## We meet your requirements

Benefit from our experience in core areas:

- Use our blood grouping monoclonal antibodies, which we supply as concentrates for further manufacturing use.
- We share our know-how in the development and production of culture media with you and produce according to your formulation.
- Our bacteriological test reagents for serotyping may be supplied as bulk reagents.

## Take advantage of our longstanding expertise in the contract manufacturing of monoclonal antibodies:

Since 1990 we have produced monoclonal antibodies for in vitro diagnostics and for research purposes. The contract manufacturing of blood-group-serological monoclonal antibodies in bulk began in 2004, in accordance with the applicable GMP regulations of the FDA.

We have a wide repertoire of processes for the cultivation, purification and labeling of monoclonal immunoglobulins

of different species (mice, rats, humans, rabbits).

From cell line entry to the finished bulk product with all necessary process stages, we are prepared to assist you with individual offers that are tailored to your specific requirements. We look forward to every new challenge.

# Contract Manufacturing

Do you have a specific request?  
Simply let us know your demand.

**sifin**

**sifin diagnostics gmbh**

Berliner Allee 317-321  
13088 Berlin, Germany

Phone: +49 30 700 144-0  
Telefax: +49 30 700 144-30  
E-Mail: [info@sifin.de](mailto:info@sifin.de)

# MICRONAUT

DIAGNOSTICS WITH PASSION



**Use the reference method for your routine diagnostics or fill the gap of your fully automated system**

**MICRONAUT systems for the identification and susceptibility testing of bacteria and yeast**

The testing principle of MICRONAUT identification systems (ID) is based on the phenotypical detection of the biochemical characteristics of microorganisms. The antimicrobial susceptibility test (AST) is based on phenotypical resistance detection by the growth of the microorganisms in the presence of the antibiotics / antifungal agents tested.

The microdilution method used here is a standardised one regarded worldwide as the recognised reference system for determining the minimum inhibitory concentration (MIC).

The substrates or antibiotics are present vacuum-dried in the microtitration plates. Thanks to a special vacuum drying procedure, these MICRONAUT plates can be stored at room temperature (15-25 °C).

Starting with a pure culture, the microorganisms are suspended in the corresponding MCN media and transferred to the MICRONAUT plates with an 8-channel pipette. Rehydration of the dried substances occurs by addition of the characterising bacteria or yeast suspension. After an incubation period of 18-24 hours the results can be photometrically measured or read visually.

As the central element in testing the MICRONAUT software combines device management, device communication and data analysis as well communication with the LIMS. The integrated expert system analyses the

photometric test measurements and assumes the clinical validation. As an option, the software can be connected bidirectionally with the laboratory EDP.



# MICRONAUT

## MICRONAUT identification

The test principle of the MICRONAUT identification systems is based on phenotypic detection of the biochemical properties of microorganisms.

Art. No.	Product	Description	Incubation time	Evaluation	Packing
M/E2-880-400	<b>MICRONAUT-GNE</b>	4 tests/plate Identification of <i>Enterobacteriaceae</i> and other gram-negative bacteria.	18 - 24 hrs	photometrically	100 plates
M/E2-710-400	<b>MICRONAUT-IDS</b>	4 tests/plate Identification of the most common clinically relevant <i>Enterobacteriaceae</i> , Non-fermenter, staphylococci, enterococci and streptococci.	5 - 6 hrs	photometrically	100 plates + 2 l NaCl
M/E5-005-200	<b>MICRONAUT-UR</b>	2 tests/plate Identification and susceptibility testing for urological diagnostics.	18 - 24 hrs	visually or photometrically	100 plates
M/E5-006-100	<b>MICRONAUT-UR-KH</b>	1 test/plate Identification and susceptibility testing for urine diagnostics, inclusive MRSA and ESBL detection.	18 - 24 hrs	photometrically	100 plates

SALE

SALE



## MICRONAUT antimicrobial susceptibility tests (AST)

The MICRONAUT AST plates are offered in individual customer defined configurations, if an agreed minimum order quantity is being purchased. The customer may choose his preferred standard like EUCAST, or CLSI and compose his own customised antibiogram from more than 200 antibiotics. All AST products listed below are available as standard products. We continuously revise the range of standard AST plates. So please contact us for details on the actual program.

### Customer defined plates

Art. No.	Product	Description	Incubation time	Evaluation	Packing
M/E1-xxx-x00	<b>MICRONAUT-S individual</b>	1 test/2 tests/4 tests/plate AST as Breakpoint or MIC method with your choice of antibiotics (minimum order quantity has to be agreed).	18 - 24 hrs	visually or photometrically	100 plates
M/EB-xxx-x00	<b>MICRONAUT-SB individual</b>	1 test/2 tests/4 tests/plate AST as Breakpoint or MIC method with your choice of antibiotics (minimum order quantity has to be agreed).	18 - 24 hrs	visually or photometrically	100 plates

### Standard plates for clinical laboratories

Art. No.	Product	Description	Incubation time	Evaluation	Packing
M/EB-422-400	<b>MICRONAUT-SB Standard Urine II (EUCAST)</b>	4 tests/plate AST as Breakpoint method for UTI relevant antibiotics, EUCAST standard.	18 - 24 hrs	visually or photometrically	100 plates
M/EB-423-200	<b>MICRONAUT-SB Varia (ambulant) (EUCAST)</b>	2 tests/plate AST as Breakpoint method for mainly ambulant relevant antibiotics, EUCAST standard.	18 - 24 hrs	visually or photometrically	100 plates

# MICRONAUT

## Standard plates for veterinary laboratories

Art. No.	Product	Description	Incubation time	Evaluation	Packing
M/E1-032-200	<b>MICRONAUT-S Mastitis</b>	2 tests/plate AST as MIC method for mastitis relevant antibiotics.	18 - 24 hrs	visually or photometrically	100 plates
M/E1-150-100	<b>MICRONAUT-S Großtiere</b>	1 test/plate AST as MIC method for antibiotics for therapy of large animals.	18 - 24 hrs	visually or photometrically	100 plates
M/E1-130-100	<b>MICRONAUT-S Kleintier</b>	1 test/plate AST as MIC method for antibiotics for therapy of small animals.	18 - 24 hrs	visually or photometrically	100 plates

## Standard plates for antifungal susceptibility testing of yeasts

Art. No.	Product	Description	Incubation time	Evaluation	Packing
M/E1-824-160	<b>MICRONAUT-AM</b>	4 tests/plate AST as Breakpoint + MIC method for antifungal agents.	22 - 48 hrs	visually or photometrically	40 plates
M/E1-831-040	<b>MICRONAUT-AM Anti Fungal Agents MIC</b>	1 test/plate AST as MIC method for antifungal agents (EUCAST).	22 - 48 hrs	visually or photometrically	40 plates
M/E1-832-080	<b>MICRONAUT-AM EUCAST AFST</b>	2 tests/plate AST as MIC method for antifungal agents (EUCAST).	22 - 48 hrs	visually or photometrically	40 plates

## MICRONAUT MIC Strips

Art. No.	Product	Description	Evaluation	Packing
M/EM-006-040	<b>MICRONAUT MIC Strip Colistin</b>	40 tests Susceptibility testing of bacteria by the broth microdilution method to determine the minimum inhibitory concentration (MIC). <u>According to the EUCAST and CLSI recommendations of March 2016 and the study of November 2016.</u>	visually	5 x 8 strips/ tests with 12 wells per strip
M/EM-024-040	<b>MICRONAUT MIC-Strip Piperacillin/Tazobactam</b>	40 tests Susceptibility testing of bacteria by the broth microdilution method to determine the MIC.	visually	5 x 8 strips/ tests with 12 wells per strip
M/EM-022-040	<b>MICRONAUT MIC Strip Vancomycin/ Teicoplanin</b>	40 tests Broth microdilution method for detection of glycopeptide resistant bacteria by determination of the MIC of Teicoplanin and Vancomycin.	visually	5 x 8 strips/ tests with 12 wells per strip



## Special plates for clinical laboratories

Art. No.	Product	Description	Incubation time	Evaluation	Packing
M/E1-111-040	<b>MICRONAUT-S β-Lactamases</b>	1 test/plate Phenotypic detection of ESBL (extended spectrum beta-lactamase), MBL (metallo-beta-lactamase), KPC (Klebsiella pneumoniae carbapenemase), AMP-C (amino-penicillin-deactivating cephalosporinase) and D-carbapenemases (OXA-48) for all relevant gram-negative bacteria in a single system.	18 - 24 hrs	visually or photometrically	40 plates
M/E1-055-040	<b>MICRONAUT-S MRSA / GP</b>	1 test/plate Detection of multidrug-resistant staphylococci (MRSA), enterococci (VRE) and pneumococci, including testing of novel antibiotics (e.g. daptomycin, ceftaroline).	18 - 24 hrs	visually or photometrically	40 plates
M/E1-114-040	<b>MICRONAUT-S MDR MRGN-Screening</b>	1 test/plate Susceptibility testing of multidrug-resistant gram-negative bacteria. Phenotypic detection of AMP-C cephalosporinase and type A-, B- and D-carbapenemases (new: ceftolozane/ tazobactam and ceftazidime/ avibactam).	18 - 24 hrs	visually or photometrically	40 plates
M/E1-085-040	<b>MICRONAUT-S Anaerobes MIC</b>	1 test/plate Susceptibility testing of treatment-relevant antimicrobial agents, such as tigecycline, moxifloxacin or ertapenem.	24 - 48 hrs	visually or photometrically	40 plates
M/E1-200-080	<b>MICRONAUT-S Anaerobier / Haemophilus (EUCAST)</b>	2 Test/Platte Susceptibility testing of anaerobe and haemophile germs.	22 - 48 hrs	visually or photometrically	40 plates
M/E1-120-080	<b>MICRONAUT-S Carbapenemases Detection</b>	2 tests/plate Phenotypic detection of clinically relevant carbapenemases OXA-48-like (type D), MBL (type B metallo-beta-lactamase) and KPC (type A Klebsiella pneumoniae carbapenemase).	18 - 24 hrs	visually or photometrically	40 plates
M/E1-973-040	<b>MICRONAUT-S Campylobacter</b>	1 test/plate Determine resistances by measuring MIC for all relevant antimicrobial agents against <i>Campylobacter</i> .	18 - 24 hrs	visually or photometrically	40 plates
M/E1-199-040	<b>MICRONAUT-S Pneumo- cocci &amp; Haemophilus MIC</b>	1 test/plate Determine resistances by measuring MIC for all relevant antimicrobial agents against pneumococci and <i>Haemophilus</i> .	18 - 24 hrs	visually or photometrically	40 plates
M/E1-099-100	<b>MICRONAUT-S Pseudomonas MIC</b>	1 test/plate Determine resistances by measuring MIC for pseudomonas-active antimicrobial agents. (new: ceftolozane/ tazobactam and ceftazidime/ avibactam).	18 - 24 hrs	visually or photometrically	100 plates

**NEW**

# MICRONAUT

## Special plates for interlaboratory comparison

Art. No.	Product	Description	Incubation time	Evaluation	Packing
M/EB-379-005	<b>MICRONAUT-SB Ringversuch Urologie</b>	1 test/plate AST towards MIC for external laboratory control for urologists.	18 - 24 hrs	visually or photometrically	5 plates
M/E1-201-005	<b>MICRONAUT-S MHK Ringversuch</b>	1 test/plate AST towards MIC for external laboratory control.	18 - 24 hrs	visually or photometrically	5 plates

## MICRONAUT reagents

Some identification tests require adding of reagent before measuring.

Art. No.	Product	Associated MCN plate	Tests	Packing
M/E2-301-001	<b>Indol Reagent</b>	MICRONAUT-IDS/UR/GNE	400	100 ml
M/E2-303-001	<b>Nitrate Reagent A</b>	Optional for MICRONAUT ID plates: MICRONAUT-IDS/UR/GNE		100 ml
M/E2-304-001	<b>Nitrate Reagent B</b>	Optional for MICRONAUT ID plates: MICRONAUT-IDS/UR/GNE		100 ml
M/E2-305-001	<b>Paraffin Oil</b>	MICRONAUT-IDS/GNE/UR	Depending on the product	100 ml
M/E2-310-001	<b>Peptidase Reagent</b>	MICRONAUT-IDS/UR	Depending on the product	100 ml
M/E2-312-001	<b>NaCl</b>	MICRONAUT identifications	Depending on the product	1 x 1000 ml
M/E2-323-001	<b>AST Reagent Kit</b>	MICRONAUT-AM		2 (2 x 4) ml

## MICRONAUT media

Bacterial suspension is prepared in MCN media.

Art. No.	Product	Associated MCN plate	Tests	Packing
M/E2-331-020	<b>Mueller Hinton Broth, cation adjusted</b>	MICRONAUT-S	20	20 x 11 ml
M/E2-331-100	<b>Mueller Hinton Broth, cation adjusted</b>	MICRONAUT-S	100	100 x 11 ml
M/E2-311-100	<b>MICRONAUT-H Medium</b>	MICRONAUT-S, fastidious microorganisms	100	100 x 11 ml
M/E2-330-020	<b>MICRONAUT-Wilkins-Chalgren Broth</b>	MICRONAUT-S Anaerob	20	20 x 11 ml
M/E2-324-020	<b>MICRONAUT-RPMI-1640 Medium + MOPS + Glucose</b>	MICRONAUT-AM	20	20 x 11 ml
M/E2-319-100	<b>MICRONAUT-SB Medium</b>	MICRONAUT-SB	100	100 x 11 ml
M/E2-338-100	<b>MICRONAUT-ID Medium</b>	MICRONAUT-UR	100	100 x 5.5 ml
M/E2-337-100	<b>MICRONAUT-AST Medium</b>	MICRONAUT-UR	100	100 x 5.5 ml
M/E2-318-200	<b>MICRONAUT-SB Medium dehydrated</b>	MICRONAUT-SB	Depending on the product used	200 g

# MICRONAUT

## MICRONAUT software

MICRONAUT software offers after automatized reading, calculation, and interpretation of identification and susceptibility testing by using MICRONAUT systems.

Art. No.	Product	Description	Packing
M/U8-305-001	<b>MICRONAUT Software</b>	Basic module	1 unit
	<b>QS Module</b>	Record of quality control data.	1 unit
	<b>Statistic Module</b>	Statistical record of resistance data.	1 unit

## Instruments

Art. No.	Product	Description	Packing
M/735391	<b>Pipette</b>	Electronic 8-channel stepper for fast inoculation of 12 x 100 µl.	1 unit
M/LT51119000	<b>Reader</b>	Reader for fast measurement of MCN plates.	1 unit
M/sifin3	<b>Reader Tecan Sunrise</b>	Reader for fast measurement of MCN plates (AST) with 3 filters.	1 unit
M/L4Y-100-001	<b>Densitometer</b>	Tube densitometer for adjustment of bacterial suspensions.	1 unit
M/2350	<b>McFarland Standard Set</b> 0.5 / 1.0 / 2.0	McFarland standards are used as a reference to adjust the density of bacterial suspensions.	3 tubes
M/709400	<b>BRAND Liquid Handling Station</b>	Liquid Handling Station (BRAND Pipetting robot) incl. software, 1 x USB interface area: 530 x 595 x 485 mm, 25 kg	1 unit
M/709426	<b>8-Channel Liquid End</b>	Pipette module for BRAND Liquid Handling Station, DE-M marking, Volume: 40-1000 µl	1 unit
M/709445	<b>Special adapter</b>	Adapter for BRAND Liquid Handling Station, for MICRONAUT reagent reservoirs, aluminium	1 unit
M/709434	<b>Tip adapter</b>	Adapter for BRAND Liquid Handling Station, for BRAND Robotic tips, aluminium	1 unit
M/709430	<b>Height adapter</b> 60 mm	Adapter for BRAND Liquid Handling Station, aluminium	1 unit
M/709432	<b>Height adapter</b> 30 mm	Adapter for BRAND Liquid Handling Station, aluminium	1 unit

## Consumables

Art. No.	Product	Description	Packing
M/ST3-001-500	<b>Matrix pipette tips</b>	For use with the Matrix pipette.	500 units
M/LH-B791204	<b>Biohit Optifit Tips Flexi-Bulk 1200</b>	For use with the Biohit pipette.	480 units
M/BH3-487-096	<b>Biohit tips rack 96</b>	For use with the Biohit pipette.	1 rack per 96 tips
M/732152	<b>BRAND Robotic tips</b>	Tips for BRAND Liquid Handling Station DE-M marking, IVD, nonsterile, PP, colourless, palletised, DNA-free, RNase-free, Endotoxin-free Volume: 40 -1000 µl Packing: 10 TipRacks	10 x 96 units
M/709458	<b>Waste Box</b>	Waste box for BRAND Liquid Handling Station holding up BRAND Robotic tips	5 units
M/R4-510-050	<b>1-Channel reservoirs</b> autoclavable	For easy plate inoculation with 1 test/plate.	50 units
M/R4-510-350	<b>1-Channel reservoirs</b> disposable product	For easy plate inoculation with 1 test/plate.	350 units
M/R4-506-050	<b>2-Channel reservoirs</b> autoclavable	For easy plate inoculation with 2 tests/plate.	50 units
M/R4-506-350	<b>2-Channel reservoirs</b> disposable product	For easy plate inoculation with 2 tests/plate.	350 units
M/R4-508-050	<b>4-Channel reservoirs</b> autoclavable	For easy plate inoculation with 4 tests/plate.	50 units
M/R4-508-350	<b>4-Channel reservoirs</b> disposable product	For easy plate inoculation with 4 tests/plate.	350 units
M/B3-002-040	<b>MICRONAUT foil</b> perforated, 40 units	For sealing of MCN identification plates (included in standard packaging).	40 units
M/B3-004-040	<b>MICRONAUT foil</b> unperforated, 40 units	For sealing of MCN AST plates (included in standard packaging).	40 units

**The MICRONAUT system prices are available on request. Please feel free to contact us.**

# Index, alphabetical

## A

<b>½ Fraser Broth Complete</b>	32
Half Fraser Broth, complete	
<b>½ Fraser Selective Supplement</b>	42
Half Fraser Selective Supplement	
<b>Acid Phosphatase Reagent</b>	44
<b>Agar for microbiological purpose</b>	46
<b>Acetamide Solution</b>	44
<b>Alkaline Peptone Water with 2% Sodium Chloride</b>	28
APW	
<b>APW</b>	28
Alkaline Peptone Water with 2% Sodium Chloride	
<b>Azide Glucose Broth</b>	28

## B

<b>Bacillus Cereus Agar (Base)</b>	28
PEMBA	
<b>Bacillus Cereus Selective Supplement</b>	42
<b>Baird Parker Agar (Base)</b>	28
<b>Bedside Cards</b>	6
<b>BHI Broth</b>	29
Brain Heart Infusion Broth	
<b>Bile Chrysoidine Glycerol Agar (Base)</b>	28
<b>Bile Chrysoidine Glycerol Agar (Base) with MUG</b>	28
<b>Bile Esculin Azide Agar</b>	28
<b>Blood Agar (Base)</b>	28
<b>BPLS Agar, modified Brilliant Green Phenol Red Agar</b>	29
acc. to Edel and Kampelmacher	
<b>BPLS-Agar acc. to Kauffmann, modified</b>	29
Brilliant Green Phenol Red Agar	
<b>BPW</b>	29
Buffered Peptone Water	
<b>Brain Heart Infusion Broth</b>	29
BHI Broth, Brain Heart Glucose Broth	
<b>Brilliant Green Bile Lactose Broth</b>	29
BRILA	
<b>Brilliant Green Phenol Red Agar</b>	29
<b>Brilliant Green Phenol Red Agar acc. to Edel and Kampelmacher</b>	29
<b>Brolacin Agar</b>	30
CLED Agar	

<b>Bromthymol Blue Broth (Base)</b>	29
<b>Buffered Nitrate Motility Medium</b>	29
<b>Buffered Peptone Water</b>	29
BPW	
<b>Buffered Sodium Chloride Peptone Solution pH 7.0</b>	29
acc. to EP/USP/JP	

## C

<b>Campylobacter Growth Supplement</b>	42
<b>Campylobacter Selective Supplement</b>	42
<b>Casein Peptone, pancreatic digested</b>	46
Tryptone	
<b>Casein Peptone Bile Agar</b>	39
Tryptone Bile Agar, TBA	
<b>Casein Soya Bean Digest Agar acc. to harm.</b>	30
EP/USP/JP CASO Agar	
<b>Casein Soya Bean Digest Broth acc. to harm.</b>	30
EP/USP/JP CASO Broth	
<b>CASO Agar</b>	30
Casein Soya Bean Digest Agar acc. to harm. EP/USP/JP	
<b>CASO Broth</b>	30
Casein Soya Bean Digest Broth acc. to harm. EP/USP/JP	
<b>CASO Broth, irradiated</b>	39
Tryptic Soy Broth, irradiated	
<b>CASO Broth, modified</b>	35
Modified Soybean Casein Digest Broth	
<b>CATC Agar (Base)</b>	30
<b>CCA</b>	30
ChroMedium Coliform, Chromogenic Coliform Agar	
<b>Cereus Selective Agar acc. to Mossel</b>	36
MYP Agar (Base)	
<b>Cetrimide Agar acc. to harm. EP/USP/JP</b>	30
<b>China Blue Lactose Agar</b>	30
<b>ChroMedium Coliform</b>	30
CCA, Chromogenic Coliform Agar	
<b>ChroMedium MLGA</b>	30
Membrane Lactose Glucuronide Agar	
<b>Chromogenic Coliform Agar</b>	30
ChroMedium Coliform, CCA	
<b>CIN Culture Medium</b>	41
Yersinia Agar (Base) acc. to Schiemann	
<b>Citrate Azide Tween Carbonate Agar</b>	30
CATC Agar (Base)	
<b>CLED Agar</b>	30

<b>Clostridial Differential Broth</b>	30
<b>Clostridial Medium acc. to harm. EP/USP/JP</b> RCM, Reinforced Medium for Clostridia acc. to harm. EP/USP/JP	38
<b>Coli Agglutinating Sera</b>	23
<b>Coliform Agar, chromogenic</b> ChroMedium Coliform, Chromogenic Coliform Agar, CCA	30
<b>Columbia Agar (Base)</b>	30
<b>Columbia Agar acc. to harm. EP/USP/JP</b>	30
<b>Columbia Blood Agar (Base)</b> Columbia Agar (Base)	30
<b>Control antigens for the Anti-Salmonella test reagents</b>	17
<b>Control antigens for the Anti-Shigella test reagents</b>	21
<b>Count Agar, sugar-free acc. to FIL-IDF</b>	30

## D

<b>DCLS Agar</b> Desoxycholate Citrate Agar modified	31
<b>Dermatophytes Agar, modified, dehydrated culture medium</b> DTM	31
<b>Desoxycholate Citrate Agar, modified</b> DCLS Agar	31
<b>DEV Gelatin Agar</b>	31
<b>DEV Lactose Peptone Broth</b>	31
<b>DEV Nutrient Agar, dehydrated culture medium</b>	31
<b>DEV Nutrient Agar, ready-to-use culture medium</b>	45
<b>Dextrose Agar</b> Glucose Agar	32
<b>DIASSALM</b> Diagnostic Semi-Solid Salmonella Agar	31
<b>Diagnostic Semi-Solid Salmonella Agar</b> DIASSALM	31
<b>Double Buffered Peptone Water</b> Peptone water, double buffered	31
<b>Double Sugar Iron Agar acc. to Kligler</b> Kligler Iron Agar, KIA	33
<b>DRCM</b> Clostridial Differential Broth	30
<b>DTM, dehydrated culture medium</b> Dermatophytes Agar, modified	31

## E

<b>Egg Yolk Emulsion</b>	42
<b>Egg Yolk Potassium Tellurite Emulsion</b>	42
<b>EMB Agar</b> Eosin Methylene Blue Lactose Saccharose Agar	32
<b>Endo Agar</b>	31
<b>Enterobacteriaceae Enrichment Broth Mossel acc. to harm. EP/USP/JP</b>	31
<b>Enterococci Selective Agar acc. to Slanetz-Bartley</b>	31
<b>Eosin Methylene Blue Lactose Saccharose Agar</b>	32

## F

<b>Fluid Thioglycolate Medium</b> Thioglycolate Medium acc. to EP/USP	39
<b>Fraser Broth (Base)</b>	32
<b>Fraser Broth, complete</b>	32
<b>½ Fraser Broth, complete</b> Half Fraser Broth, complete	32
<b>Fraser Selective Supplement</b>	42
<b>½ Fraser Selective Supplement</b>	42

## G

<b>Gassner Agar, modified</b> Water-blue Metachrome yellow Lactose Agar acc. to Gassner	41
<b>GCG Agar</b> Bile Chrysoidine Glycerol Agar (Base)	28
<b>Gelatine Lactose Medium</b>	32
<b>Glucose Agar</b>	32
<b>Glucose for microbiological use</b>	46
<b>Glucose Nutrient Broth</b>	32
<b>Glucose Yeast Extract Cysteine Agar (Base) acc. to Beerens</b>	32
<b>Glycerin, water-free</b>	44
<b>GSP Agar (Base)</b>	32

# Index, alphabetical

## H

<b>Half Fraser Broth, complete</b>	32
<b>Half Fraser Selective Supplement</b>	42
<b>HIB Medium (Base)</b>	33

## I

<b>Iron Double Sugar Agar acc. to Kligler KIA</b> Kligler Iron Agar, Double Sugar Iron Agar acc. to Kligler	33
--	----

## K

<b>Kanamycin Esculin Azide Agar</b> KAA Agar	33
<b>Kanamycin Esculin Azide Broth</b> KAA Broth	33
<b>KIA</b> Kligler Iron Agar	33
<b>King's Agar (Base)</b>	33
<b>Kligler Iron Agar</b> Double Sugar Iron Agar acc. to Kligler, KIA	33

## L

<b>Lactobacillus Agar acc. to de Man, Rogosa and Sharpe</b> MRS Agar (Base)	35
<b>Lactose Broth</b>	33
<b>Lactose for microbiology</b>	46
<b>Lactose Fuchsin Sulfite Agar</b> Endo Agar	31
<b>Lactose TTC Agar with Tergitol-7</b>	33
<b>Lauryl Sulfate Broth</b> Lauryl Sulfate Tryptose Lactose Broth	33
<b>Lauryl Sulfate Tryptose Broth with Tryptophan and MUG</b>	33
<b>Lauryl Sulfate Tryptose Lactose Broth</b> Lauryl Sulfate Broth	33
<b>LB Agar</b> Luria Bertani Agar acc. to Miller	33
<b>LB Broth</b> Luria Bertani Medium acc. to Miller	34
<b>LDC Broth</b> Lysine Decarboxylase Broth	34
<b>Leifson Agar, modified</b> Desoxycholate Citrate Agar (DCLS Agar), modified	31

<b>Linden Grain Medium</b>	34
<b>LST/MUG Medium</b> Lauryl Sulfate Tryptose Broth with Tryptophan and MUG	33
<b>Luria Bertani Agar acc. to Miller</b> LB Agar	33
<b>Luria Bertani Medium acc. to Miller</b> LB Broth	34
<b>Lysine Agar, ready-to-use culture medium</b>	45
<b>Lysine Decarboxylase Broth</b> LDC Broth	34

## M

<b>MacConkey Agar</b>	34
<b>MacConkey Agar acc. to harm. EP/USP/JP</b>	34
<b>MacConkey Broth acc. to harm. EP/USP/JP</b>	34
<b>Magnesium Chloride Malachite Green Broth acc. to Rappaport-Vassiliadis</b>	34
<b>Malachite-green Broth</b>	34
<b>Malt Extract Agar</b>	34
<b>Mannitol Egg Yolk Polymyxin Agar</b> MYP Agar (Base)	35
<b>Mannitol Salt Agar acc. to harm. EP/USP/JP</b>	35
<b>Maximum Recovery Diluent</b> Peptone Saline Diluent	37
<b>m-CP Agar (Base)</b>	35
<b>m-CP Selective Supplement</b>	43
<b>Meat extract</b>	46
<b>Meat peptone, pancreatic digested</b>	46
<b>MegaPec Broth, ready-to-use culture medium</b>	45
<b>Membrane Lactose Glucuronide Agar</b> ChroMedium MLGA	30
<b>m-Enterococci Agar</b> Enterococci Selective Agar acc. to Slanetz-Bartley	31
<b>Methyl-red Voges Proskauer Broth</b> MR-VP	35
<b>MICRONAUT system</b>	53
<b>Minca-Agar, modified, dehydrated culture medium</b>	35
<b>Minca Supplement</b>	43
<b>MLGA (ChroMedium MLGA)</b> Membrane Lactose Glucuronide Agar	30



<b>Modified Scholtens' Broth</b>	35
<b>Modified Semi-solid Rappaport Vassiliadis Medium</b> MSRV Medium (Base)	35
<b>Modified Soybean Casein Digest Broth</b> CASO Broth, modified	35
<b>Monoclonal antibodies</b>	49
<b>MRS Agar (Base)</b>	35
<b>MRS Agar (Base) with pH 5.7</b>	35
<b>MRS Broth (Base)</b>	35
<b>MSB</b> Modified Scholtens' Broth	35
<b>MSRV Medium (Base)</b>	35
<b>Mueller Hinton Agar</b>	35
<b>Mueller Hinton Broth</b>	36
<b>MYP Agar (Base)</b>	36

## N

<b>Nessler's Reagent</b>	44
<b>Novobiocin Selective Supplement</b>	43
<b>Nutrient Agar I</b>	36
<b>Nutrient Agar III</b>	36
<b>Nutrient Agar acc. to German drinking water regulation and Food and Feed Code</b>	36
<b>Nutrient Broth</b>	36
<b>Nutrient Broth I</b>	36
<b>Nutrient Broth II</b>	36

## O

<b>Orange Serum Agar</b>	36
<b>Orange Serum Broth</b>	37
<b>Oxidase Reagent</b>	44

## P

<b>Palcam Agar (Base)</b> Palcam Agar acc. to van Netten	37
<b>Palcam Selective Supplement</b>	43
<b>Peptone Saline Diluent</b> Maximum Recovery Diluent	37
<b>Peptone Water, Buffered</b> Buffered Peptone Water	29
<b>Peptone Water, Double Buffered</b> Double Buffered Peptone Water	31
<b>Peptone Water with 2% Sodium Chloride, alkaline</b> Sodium Chloride Polymyxin Solution (Base)	28
<b>PK® Products</b>	8
<b>Plate Count Agar</b>	37
<b>Plate Count Agar with Skimmed Milk</b>	37
<b>Polymyxin B Selective Supplement</b> Bacillus Cereus Selective Supplement	42
<b>Polyoxyethylene sorbitan monooleate</b> Tween® 80	44
<b>Potato Dextrose Agar acc. to harm. EP/USP/JP</b>	37
<b>Pseudomonas Aeromonas Agar acc. to Kielwein</b> GSP Agar (Base)	32
<b>Pseudomonas Agar (Base)</b>	37
<b>Pseudomonas CN Selective Supplement</b>	43

## R

<b>R2A Agar</b>	37
<b>Rabies Detection Product</b>	47
<b>Rappaport Vassiliadis Salmonella Enrichment Broth acc. to harm. EP/USP/JP</b> RVS Broth	37
<b>RCM acc. to harm. EP/USP/JP</b> Reinforced Medium for Clostridia acc. to harm. EP/USP/JP	38
<b>Recovery Diluent, maximum</b> Peptone Saline Diluent	37
<b>Reinforced Medium for Clostridia acc. to harm. EP/USP/JP</b> RCM acc. to harm. EP/USP/JP	38
<b>Rice Extract Agar</b>	38
<b>RV Medium</b> Magnesium Chloride Malachite Green Broth acc. to Rappaport-Vassiliadis	34
<b>RVS Broth acc. to harm. EP/USP/JP</b> Rappaport Vassiliadis Salmonella Enrichment Broth acc. to harm. EP/USP/JP	37

# Index, alphabetical

## S

<b>Sabouraud Dextrose Agar</b>	38
Sabouraud 2% Glucose Agar	
<b>Sabouraud 2%Glucose Agar</b>	38
<b>Sabouraud Dextrose Agar acc. to harm. EP/USP/JP</b>	38
<b>Sabouraud Dextrose Broth acc. to harm. EP/USP/JP</b>	38
<b>Sabouraud 2%Glucose Chloramphenicol Agar</b>	38
<b>Sabouraud Glucose Medium with Chloramphenicol</b>	38
<b>Sabouraud Glucose Slant Agar, ready-to-use culture medium</b>	45
<b>Sabouraud Glucose Slant Agar with Chloramphenicol and Cycloheximide, ready-to-use culture medium</b>	45
<b>Salmonella Agglutinating Sera</b>	12
<b>Salmonella Shigella Agar, modified</b>	39
SS Agar, modified	
<b>Schaedler Agar (Base)</b>	38
<b>Schaedler Broth</b>	38
<b>Scholtens' Broth modified</b>	35
Modified Scholtens' Broth	
<b>Selenite Broth</b>	38
<b>Selenite Cystine Broth</b>	38
<b>Serafol</b>	6
<b>Shigella Agglutinating Sera</b>	20
<b>Simmons Citrate Agar</b>	38
<b>SMAC Agar</b>	39
Sorbitol MacConkey Agar	
<b>Sodium chloride for microbiological purpose</b>	46
<b>Sodium Chloride Peptone Solution pH 7.0 acc. to EP/USP/JP, buffered</b>	29
<b>Sodium Chloride Polymyxin Solution (Base)</b>	28
Alkaline Peptone Water with 2% Sodium Chloride, APW	
<b>Sorbitol MacConkey Agar</b>	39
SMAC Agar	
<b>Soyabean Casein Digest Broth, modified</b> Modified	35
Soybean Casein Digest Broth, CASO Broth modified	
<b>Soyabean Casein Digest Broth, irradiated</b>	39
Tryptic Soy Broth irradiated, CASO Broth irradiated	
<b>Soy bean Peptone, papainic digested</b>	46
<b>SS Agar, modified</b>	39
Salmonella Shigella Agar, modified	
<b>Swarm Agar, ready-to-use culture medium</b>	45

<b>Synthetic Nutrient Deficient Agar</b>	39
SNA	

## T

<b>TBA</b>	39
Tryptone Bile Agar	
<b>TBX Chromogen Agar</b>	39
<b>Tergitol 7 TTC Agar</b>	33
Lactose TTC Agar with Tergitol-7	
<b>Tetrathionate Broth (Base) acc. to Muller-Kauffmann</b>	39
<b>TGE Agar</b>	38
Plate Count Agar	
<b>Thioglycolate Medium acc. to EP/USP</b>	39
<b>Thioglycolate Medium, modified</b>	39
<b>Thioglycolate Medium without Indicator</b>	39
<b>Tryptic Soy Broth (CASO), irradiated</b>	39
<b>Tryptone</b>	46
Casein peptone, pancreatic digested	
<b>Tryptone Bile Agar</b>	39
TBA	
<b>Tryptone Bile X Glucuronid Agar</b>	39
TBX Chromogen Agar	
<b>Tryptone Glucose Yeast Extract Agar</b>	37
Plate Count Agar	
<b>Tryptone Glucose Yeast Extract Milk Agar</b>	37
Plate Count Agar with Skimmed Milk	
<b>Tryptone Soy Yeast Extract Agar</b>	40
TSYEA	
<b>Tryptophan Broth</b>	40
Tryptophan Pepton Water	
<b>Tryptophan Pepton Water</b>	40
<b>Tryptose Sulfit Cycloserine Agar</b>	40
TSC Agar (Base)	
<b>TSC Agar (Base)</b>	40
Tryptose Sulfit Cycloserine Agar	
<b>TSC Selective Supplement</b>	43
<b>TSYEA</b>	40
Tryptone Soy Yeast Extract Agar	
<b>Tween® 80</b>	44

## U

<b>Urea Agar (Base) acc. to Christensen</b>	40
<b>Urea Indole Motility Medium</b>	33
HIB Medium (Base)	
<b>Urea Solution, 40%</b>	44

## V

<b>VLB-S7-S Agar, ready-to-use culture medium</b>	45
<b>VLB-S7-S Broth, ready-to-use culture medium</b>	45
<b>Violet Red Bile Glucose Agar</b>	40
VRBD Agar	
<b>Violet Red Bile Glucose Agar acc. to harm. EP/USP/JP</b>	40
VRBD Agar acc. to harm. EP/USP/JP	
<b>Violet Red Bile Lactose Agar</b>	40
<b>VRB Agar</b>	40
Violet Red Bile Lactose Agar	
<b>VRBD Agar</b>	40
Violet Red Bile Glucose Agar	
<b>VRBD Agar acc. to harm. EP/USP/JP</b>	40
Violet Red Bile Glucose Agar acc. to harm. EP/USP/JP	

## W

<b>Water-blue Metachrome-yellow Lacose Agar acc. to Gassner, modified</b>	41
<b>WLD, ready-to-use culture medium</b>	45
WL Nutrient Agar with Cycloheximide	
<b>WL Nutrient Agar, ready-to-use culture medium</b>	45
WLN	
<b>WL Nutrient Agar with Cycloheximide, ready-to-use culture medium</b> WLD	45
<b>WLN, ready-to-use culture medium</b>	45
WL Nutrient Agar	
<b>Wort Agar (Base)</b>	41
<b>Wort Agar, ready-to-use culture medium</b>	45
<b>Wort Broth (Base)</b>	41
<b>Wort Peptone Agar</b>	34
Malt Extract Agar	

## X

<b>XLD Agar acc. to ISO 6579</b>	41
<b>XLD Agar acc. to harm. EP/USP/JP</b>	41

<b>Xylose Lysine Desoxycholate Agar</b>	41
XLD Agar acc. to ISO 6579	

<b>Xylose Lysine Desoxycholate Agar acc. to harm. EP/USP/JP</b>	41
XLD Agar acc. to harm. EP/USP/JP	

## Y

<b>Yeast Extract Agar</b>	41
---------------------------	----

<b>Yeast extract for microbiological use</b>	46
--	----

<b>Yeast Extract Glucose Chloramphenicol Agar</b>	41
YGC Agar	

<b>Yersinia Agar (Base) acc. to Schiemann</b>	41
---	----

<b>Yersinia Agglutinating Sera</b>	22
------------------------------------	----

<b>Yersinia Selective Supplement</b>	43
--------------------------------------	----

<b>YGC Agar</b>	41
Yeast Extract Glucose Chloramphenicol Agar	

# Certificates

Translation from German



## CERTIFICATE

The certification body confirms to

**sifin diagnostics gmbh**  
**Berliner Allee 317-321**  
**13088 Berlin**  
**Germany**

for the development, manufacturing and sale of products for human and veterinary medical in-vitro-diagnostics as well as for the microbiological examination of water and food and other diagnostic applications the conformity of the introduced quality management system with the standard

### DIN EN ISO 9001:2015

Start of validity:	07.07.2017
End of validity:	06.07.2020
Report and certificate number:	IC00016 038 17
The certificate consists of	1 page

*This certificate includes an annual examination of the QMS by IFTA AG, according to the specified standard.*

Berlin, 06.07.2017



Prof. Dr. Jörn Karge  
CEO



IFTA AG | Neukirchstraße 26 | 13089 Berlin | fon 0049.30.47 88 03 0 | fax 0049.30.47 88 03 20 | web www.ifta-ag.de  
Rev.: 1

# Certificate

**mdc medical device certification GmbH**  
certifies that

## sifin

**sifin diagnostics gmbh**  
**Berliner Allee 317-321**  
**13088 Berlin**  
**Germany**

for the scope

**development, manufacturing and distribution of  
in vitro diagnostic medical devices for the product groups:  
blood grouping, bacteriological test reagents and culture media as well as  
manufacturing of raw materials for manufacturing of  
in vitro diagnostic medical devices**

has introduced and applies a

## Quality Management System

The mdc audit has proven that this quality management system  
meets all requirements of the following standard

### EN ISO 13485

Medical devices – Quality management systems –  
Requirements for regulatory purposes

EN ISO 13485:2016 + AC:2016 - ISO 13485:2016

Valid from	2018-10-23
Valid until	2021-10-22
Registration no.	D1058700042
Report no.	P18-00745-121758
Stuttgart	2018-07-16

  
Head of Certification Body



mdc medical device certification GmbH  
Kriegerstraße 6  
D-70191 Stuttgart, Germany  
Phone: +49-(0)711-253597-0  
Fax: +49-(0)711-253597-10  
Internet: <http://www.mdc-ce.de>

For electronic publication only

# Certificates



Deutsche Akkreditierungsstelle GmbH

Annex to the Accreditation Certificate D-PL-20980-01-00  
according to DIN EN ISO/IEC 17025:2005

Period of validity: 12.07.2018 to 11.07.2023      Date of issue: 12.07.2018

Holder of certificate:

**sifin diagnostics gmbh**  
**Prüflaboratorium Nährmedien**  
**Berliner Allee 317-321, 13088 Berlin**

Tests in the fields:

**microbiological analysis of culture media at area food, animal feed and water**

**The testing laboratory is permitted, without being required to inform and obtain prior approval from DAkkS, to use standards or equivalent testing methods listed here with different issue dates.**

DIN EN ISO 11133 2015-01	Microbiology of food, animal feed and water - Preparation, production, storage and performance testing of culture media ( <i>withdrawn standard</i> )
-----------------------------	---

**Abbreviations used:**

DIN	German Institute for Standardisation (Deutsches Institut für Normung e. V.)
EN	European Standard
IEC	International Electrotechnical Commission
ISO	International Organization for Standardization

# Distributors

## Albania

### Bacteriological test reagents

Krijon Ltd.  
Phone: +355-4-2347713  
Fax: +355-4-2340817  
E-Mail: krijon@albmail.com

## Austria

### Bacteriological test reagents

#### Culture media

MICRONAUT  
BIOMEDICA  
Medizinprodukte GmbH & Co KG  
(DiaMondial Group)  
Phone: +43-1-291-07-56  
Fax: +43-1-290-14-29  
E-Mail: sales.biomedica@bmgrp.at  
www.biomedica.co.at

## Belarus

### Bacteriological test reagents

Himhrom  
Phone: +375-172-161797  
Fax: +375-172-161798  
E-Mail: himhrom@open.by

## Bulgaria

### Bacteriological test reagents

MICRONAUT  
Ridacom Ltd.  
Phone: +359-2-9559998  
E-Mail: info@ridacom.com  
www.ridacom.com

### Blood group products

IVD Bulgaria  
Phone: +359-2-49-11-226  
Fax: +359-2-97-58-023  
E-Mail: ivd@ivd.bg  
www.ivd.bg

## MICRONAUT

AQUACHIM JSco.  
Phone: +359-2-807-50-25  
Fax: +359-2-807-50-50  
E-Mail: aquachim@aquachim.bg  
www.aquachim.bg

## Chile

### Bacteriological test reagents

### Blood group products

Farmalatina Ltda  
Phone: +56-2-838-5000  
Fax: +56-2-838-5050  
E-Mail: farmalatina@farmalatina.cl  
www.farmalatina.cl

## Croatia

### Bacteriological test reagents

Labomar d.o.o.  
Phone: +385-1-3457969  
Fax: +385-1-3457967  
E-Mail: viktor@labomar.hr  
www.labomar.hr

## Cyprus

### Bacteriological test reagents

#### Culture media

Scientronics Ltd.  
Phone: +357-22-467880  
Fax: +357-22-764614  
E-Mail: a.sarpetsas@biotronics.com.cy  
www.scientronics.com

## Czech Republic

### Bacteriological test reagents

BioVendor – Laboratorní medicína a.s.  
Phone: +420-549-124-111  
Fax: +420-549-211-465  
E-Mail: info@biovendor.cz  
www.biovendor.cz

### Bacteriological test reagents

Eurex Medica, spol. s.r.o.  
Phone: +420-599-526-510  
Fax: +420-596-614-507  
E-Mail: info@eurexmedica.cz  
www.eurexmedica.cz

## Egypt

### Bacteriological test reagents

Sigma Trade Co.  
Phone: +20-2-5737861  
Fax: +20-2-5737861  
E-Mail: info@sigmatrade.net  
www.sigmatrade.net

## France

### Bacteriological test reagents

Eurobio (DiaMondial Group)  
Phone: +33-1-69-07-94-77  
Fax: +33-1-69-07-95-34  
E-Mail: adv@eurobio.fr  
www.eurobio.fr

## MICRONAUT

SOBioda  
Phone: +33-4-76610202  
Fax: +33-4-76619181  
E-Mail: info@sobioda.com  
www.sobioda.eu

## Georgia

### Bacteriological test reagents

Human Diagnostic Georgia  
Phone: +995-32-295-02-22  
Fax: +995-32-296-28-60  
E-Mail: humangeorgia@caucasus.net

## Greece

### Blood group products

Vaktro Scientific  
Phone: +30-2610-223999  
Fax: +30-2610-223595  
E-Mail: info@vaktro.gr  
www.vaktro.gr

### Blood group products

Biosna Basklavanis - Fotoglou - Perros S.A.  
Phone: +30-210-7799468  
Fax: +30-210-7791439  
E-Mail: biosna@otenet.gr

## MICRONAUT

Bio-G  
Phone: +30-2310-558888  
Fax: +30-2310-558888  
E-Mail: info@biog.gr  
www.biog.gr

## Hungary

### Bacteriological test reagents

#### Culture media

### Blood group products

Biotest Hungaria Kft.  
Phone: +36-23-511311  
Fax: +36-23-511310  
E-Mail: biotest@biotest.hu  
www.biotest.hu

## Ireland

### Bacteriological test reagents

Cruinn Diagnostics Ltd.  
(DiaMondial Group)  
Phone: +353-1-6297400  
Fax: +353-1-6297401  
E-Mail: info@cruinn.ie  
www.cruinn.ie

## Italy

### Bacteriological test reagents

#### Culture media

Generon S.p.A.  
Phone: +39-059-8637161  
Fax: +39-059-7353024  
E-Mail: marketing@generon.it  
www.generon.it

### Bacteriological test reagents

D.I.D S.p.a (DiaMondial Group)  
Phone: +39-02-40-09-02-22  
Fax: +39-02-48-70-58-02  
E-Mail: info@did.it  
www.did.it

## Kosovo

### Bacteriological test reagents

#### Culture media

ERAMED  
Phone: +381-38-245-361  
Fax: +381-38-245-361  
E-Mail: flamur@era-med.com  
www.era-med.com

## Latvia

### Bacteriological test reagents

#### Culture media

Enola SIA  
Phone: +371-6-7372566  
Fax: +371-6-7379302  
E-Mail: info@enola.lv  
www.enola.lv

## Lebanon

### Blood group products

BioCare  
Phone: +961-1-686033  
Fax: +961-1-686032  
E-Mail: biocare@biocarelb.com  
www.biocarelb.com

## Lithuania

### Bacteriological test reagents

Biometrija UAB  
Phone: +370-5-2481824  
Fax: +370-5-2400085  
E-Mail: biometrija@biometrija.lt  
www.biometrija.lt

## MICRONAUT

UAB „Interautomatika“  
Phone: +370-5-2607810  
Fax: +370-5-2411464  
E-Mail: alla@interlab.lt  
www.interautomatika.lt

## Macedonia

### Bacteriological test reagents

### Blood group products

Biotek doo Skopje  
Phone: +389-23-110-106  
Fax: +389-23-126-274  
E-Mail: info@biotek.com.mk  
www.biotek.com.mk

## Malaysia

## MICRONAUT

LABLINK  
Phone: +60-3-40234588  
Fax: +60-3-40234298  
E-Mail: purchasing@lablink.com.my  
www.lablink.com.my

## Malta

### Bacteriological test reagents

#### Culture media

E.J. Busuttill Ltd.  
Phone: +356-21-447184  
Fax: +356-21-445366  
E-Mail: info@ejbusuttill.com  
www.ejbusuttill.com

## Moldova

### Bacteriological test reagents

IM Global Biomarketing Group  
Phone: +373-22-54-73-73  
Fax: +373-22-54-73-73  
E-Mail: office@gbg.md  
www.gbg.md

## Mongolia

## MICRONAUT

Monbiopharm Co. Ltd.  
Phone: +976-99-156493  
Fax: +976-11-324420  
E-Mail: delgermaa.e@monbiopharm.mn  
www.monbiopharm.mn

## Montenegro

### Bacteriological test reagents

Morazzia Laboratory  
Phone: +382-40-235500  
Fax: +382-40-257035  
E-Mail: morazzia@cg.yu



## Morocco

### Bacteriological test reagents

#### Blood group products

Top Labo  
Phone: +212- 52-2989221  
Fax: +212-52-2989223  
E-Mail: toplabo0@gmail.com

## Netherlands

### Bacteriological test reagents

#### Culture media

#### MICRONAUT

bioTRADING Benelux B. V.  
(DiaMondial Group)  
Phone: +31-297-286848  
Fax: +31-297-287570  
E-Mail: info@biotrading.com  
www.biotrading.com

## New Zealand

### Bacteriological test reagents

Ngaio Diagnostics  
Phone: +64-3-5484727  
Fax: +64-3-5484729  
E-Mail: info@ngaio.co.nz  
www.ngaio.co.nz

## Norway

### Bacteriological test reagents

#### MICRONAUT

Montebello Diagnostics AS  
Phone: +47-22141490  
Fax: +47-22144490  
E-Mail: mail@montebello.no  
www.montebello.no

## Panama

### Bacteriological test reagents

Inversiones Sagrav S.A.  
Phone: +507-2330902  
Fax: +507-2335685  
E-Mail: sagrav@sagrav.com.pa

## Philippines

### Bacteriological test reagents

Goldquest Biotechnologies Inc.  
Phone: +63-2-2419515  
Fax: +63-2-2433486  
E-Mail: quest-diagnostic@pldtdsl.net  
www.goldquest-biotechnologies.com

#### MICRONAUT

Scientific Biotech Specialties Inc  
Phone: +63-2-8996557  
Fax: +63-2-8969382  
E-Mail: sbsi.diag@sbsi.com.ph  
www.sbsi.com.ph

## Poland

### Bacteriological test reagents

Nobipharm Sp.z.o.o.  
Phone: +48-22-6339802  
Fax: +48-22-6339802  
E-Mail: info@nobipharm.pol.pl  
www.nobipharm.pol.pl

## Bacteriological test reagents

Graso Zenon Sobiecki  
Phone: +48-58-562-30-21  
Fax: +48-58-562-79-87  
E-Mail: zamowienia@graso.com.pl  
www.podloza.pl

## Portugal

### Blood group products

Iberlab & Immunoreage solucoes  
para laboratório, lda.  
Phone: +351-22-2087876  
Fax: +351-22-2054328  
E-Mail: iberlab@iberlab.pt  
www.iberlab.pt

#### MICRONAUT

Quilaban  
Phone: +351-21-9236372  
Fax: +351-21-9236389  
E-Mail: k.ferreira@quilaban.pt  
www.quilaban.pt

## Romania

### Culture media

SC EMSAR SRL  
Phone: +40-31-1013182  
Fax: +40-21-6671243  
E-Mail: office@emsar.ro  
www.emsar.ro

### Bacteriological test reagents

Eurovet Lab  
Phone: +40-21-3102153  
Fax: +40-21-3102154  
E-Mail: cosmin@eurovet-lab.ro

### Blood group products

SC Vitro Biochem srl  
Phone: +40-21-4118833  
Fax: +40-21-4116023  
E-Mail: office@vitro.ro

## Russia

### Bacteriological test reagents

#### Culture media

HEM Ltd.  
Phone: +74-95-7870432  
Fax: +74-95-6120481  
E-Mail: sale@hemltd.ru  
www.hemltd.ru

#### MICRONAUT

SY-LAB ooo  
Phone: +495 739 58 55  
Fax: +495 739 58 56  
E-Mail: jane@sy-lab.ru  
www.sy-lab.ru

## Saudi-Arabia

### Bacteriological test reagents

Gamsco  
Phone: +966-3-839-4085  
Fax: +966-3-839-4084  
E-Mail: sales@gamscomed.com  
www.gamscomed.com

## Slovenia

### Bacteriological test reagents

Mediline d.o.o.  
Phone: +386-1-8308040  
Fax: +386-1-8308070  
E-Mail: info@mediline.si  
www.mediline.si

## South Africa

### Blood group products

Ethitech  
Phone: +27-21-855-0307  
Fax: +27-86-657-1986  
E-Mail: sarah@ethitech.co.za  
www.ethitech.co.za

## Spain

### Bacteriological test reagents

#### Blood group products

Werfen Iberia  
(DiaMondial Group)  
Phone: +34-934-010211  
Fax: +34-934-010350  
E-Mail: allenderrozos@werfen.com  
www.werfen.com

## Sweden

#### MICRONAUT

Miclev AB  
Phone: +46-40-365400  
E-Mail: elin.wallin@miclev.se  
www.miclev.se

## Switzerland

### Bacteriological test reagents

#### MICRONAUT

Biotest (Schweiz) AG  
Phone: +41-62-8890000  
Fax: +41-62-8890001  
E-Mail: mail.ch@biotest.com  
www.biotest.ch

## Taiwan

### Bacteriological test reagents

Linkseas Trading Co., Ltd.  
Phone: +886-2-27735142  
Fax: +886-2-27721983  
E-Mail: lks1976.tw@msa.hinet.net

## Turkey

### Blood group products

Toprak Medikal Ltd.  
Phone: +90-212-6217016  
Fax: +90-212-6217018  
E-Mail: topramed@superonline.com

## Ukraine

### Bacteriological test reagents

NPF Ukrmedialab Ltd.  
Fon: +380-667-207-941  
E-Mail: himediaua@gmail.com

## Bacteriological test reagents

### Culture media

#### MICRONAUT

„SORS UKRAINE“ LLC  
Phone: +384-446-76441  
Fax: +384-446-76441  
E-Mail: ierokhina@sors.com.ua

## United Arab Emirates

#### MICRONAUT

Alliance Global Group  
Phone: +971 (4) 375-27-44  
Fax: +971 (4) 429-13-76  
E-Mail: customerservice@agbl.net

## United Kingdom

### Bacteriological test reagents

#### MICRONAUT

BioConnections  
Phone: +44-1782-516010  
Fax: +44-1782-510733  
E-Mail: inbox@bcnx.co.uk  
www.bioconnections.co.uk

## Venezuela

### Bacteriological test reagents

#### Blood group products

#### Culture media

Ganbaro  
Phone: +58-212-5410501  
Fax: +58-212-5410866  
E-Mail: gerencia.producto@ganbaro.com.ve  
www.ganbaro.com.ve

## Vietnam

### Blood group products

DEKA Co., Ltd., North Vietnam  
Phone: +84-4-37830363  
Fax: +84-4-37830364  
E-Mail: deka@hn.vnn.vn  
www.deka.vn

### Blood group products

Phuong Dong TECHME Co., Ltd.  
Phone: +84-4-66872909  
E-Mail: vinhtruongtby@gmail.com

## Vietnam

### Blood group products

VAN XUAN, South Vietnam  
Phone: +84-8-38-686-109  
Fax: +84-8-38-686-157  
E-Mail: triphuong@hcm.fpt.vn

### Blood group products

DEKA JOINT STOCK CO., South Vietnam  
Phone: +84-8-62-638-402  
Fax: +84-8-62-638-451  
E-Mail: dekahcm@hcm.vnn.vn  
www.dekavn.com

#### MICRONAUT

Eastern Medical Equipment  
Co.Ltd., Hanoi  
Phone: +84-983-184-161  
E-Mail: thuy.nt\_hn@eastern.vn  
www.easterngroup.com.vn



Do you have a specific request?  
Simply let us know your demand.

sifin

sifin diagnostics gmbh

Berliner Allee 317-321  
13088 Berlin, Germany

Phone: +49 30 700 144-0  
Telefax: +49 30 700 144-30  
E-Mail: [info@sifin.de](mailto:info@sifin.de)



**EUREX** *Medica*®

**EUREX MEDICA, spol. s r.o.**

Výstavní 604/111

703 00 Ostrava – Vítkovice

Tel.: 599 526 510,

Fax: 596 614 507

e-mail: [expedice@eurexmedica.cz](mailto:expedice@eurexmedica.cz)

web: <http://www.eurexmedica.cz>

# sifin

sifin diagnostics gmbh

Berliner Allee 317-321  
13088 Berlin, Germany

Phone: +49 30 700 144-0  
Telefax: +49 30 700 144-30  
E-Mail: [info@sifin.de](mailto:info@sifin.de)



**EUREX** *Medica*®

**EUREX MEDICA, spol. s r.o.**

Výstavní 604/111

703 00 Ostrava – Vítkovice

Tel.: 599 526 510,

Fax: 596 614 507

e-mail: [expedice@eurexmedica.cz](mailto:expedice@eurexmedica.cz)

web: <http://www.eurexmedica.cz>

[www.sifin.de](http://www.sifin.de)